arXiv:math/0609741v7 [math.AG] 15 Feb 2010

Poisson deformations of affine symplectic varieties

Yoshinori Namikawa

Introduction

A symplectic variety X is a normal algebraic variety (defined over $\mathbf{C}$) which admits an everywhere non-degenerate d-closed 2-form ω on the regular locus X_{reg} of X such that, for any resolution $f : \tilde{X} \to X$ with $f^{-1}(X_{reg}) \cong X_{reg}$, the 2-form ω extends to a regular closed 2-form on $\tilde{X}$ (cf. [Be]). There is a natural Poisson structure $\{\ ,\ \}$ on X determined by ω. Then we can introduce the notion of a Poisson deformation of $(X, \{\ ,\ \})$. A Poisson deformation is a deformation of the pair of X itself and the Poisson structure on it. When X is not a compact variety, the usual deformation theory does not work in general because the tangent object $\mathbf{T}^1_X$ may possibly have infinite dimension, and moreover, infinitesimal or formal deformations do not capture actual deformations of non-compact varieties. On the other hand, Poisson deformations work very well in many important cases where X is not a complete variety. Denote by PD_X the Poisson deformation functor of a symplectic variety (cf. §1). In this paper, we shall study the Poisson deformation of an affine symplectic variety. The main result is:

Theorem (5.1). *Let X be an affine symplectic variety. Then the Poisson deformation functor* PD_X *is unobstructed.*

A Poisson deformation of X is controlled by the Poisson cohomology $\mathrm{HP}^2(X)$ (cf. [G-K], [Na 2]). When X has only terminal singularities, we have $\mathrm{HP}^2(X) \cong H^2((X_{reg})^{an}, \mathbf{C})$, where $(X_{reg})^{an}$ is the complex space associated with X_{reg}. In that case this description enables us to prove that PD_X is unobstructed ([Na 2], Corollary 15). But, in general, there is no such direct, topological description of $\mathrm{HP}^2(X)$. Let us explain our strategy to describe $\mathrm{HP}^2(X)$. As remarked, $\mathrm{HP}^2(X)$ is identified with $\mathrm{PD}_X(\mathbf{C}[\epsilon])$ where $\mathbf{C}[\epsilon]$ is the ring of dual numbers over $\mathbf{C}$. First, note that there is an open locus U of X where X is smooth, or is locally a trivial deformation of a (surface) rational double point at each $p \in U$. Let Σ be the singular locus of U. Note that $X \setminus U$ has codimension ≥ 4 in X (cf. [Ka 1]). Moreover, we have $\mathrm{PD}_X(\mathbf{C}[\epsilon]) \cong \mathrm{PD}_U(\mathbf{C}[\epsilon])$. Put $T^1_{U^{an}} := \underline{\mathrm{Ext}}^1(\Omega^1_{U^{an}}, \mathcal{O}_{U^{an}})$. As is well-known, a (local) section of $T^1_{U^{an}}$ corresponds to a 1-st order deformation of U^{an}. In §1, we shall construct a locally constant sheaf $\mathcal{H}$ of $\mathbf{C}$-modules as a subsheaf of $T^1_{U^{an}}$. The sheaf $\mathcal{H}$ is intrinsically characterized as the sheaf of germs of sections of $T^1_{U^{an}}$ *which come from Poisson deformations of* U^{an} (cf. Lemma (1.5)). Now we have an exact sequence (cf. Proposition (1.11)):

$$0 \to H^2(U^{an}, \mathbf{C}) \to \mathrm{PD}_U(\mathbf{C}[\epsilon]) \to H^0(\Sigma, \mathcal{H}).$$

Here the first term $H^2(U^{an}, \mathbf{C})$ is the space of locally trivial[1] Poisson deformations of U. By the definition of U, there exists a minimal resolution $\pi : \tilde{U} \to U$. Let m be the number of irreducible components of the exceptional divisor of π. Section 3 is a preliminary section for section 4. However, Proposition (3.2) is the core of the argument in §4. The main result of §4 is:

Proposition (4.2). *The following equality holds:*

$$\dim H^0(\Sigma, \mathcal{H}) = m.$$

In order to prove Proposition (4.2), we need to know the monodromy action of $\pi_1(\Sigma)$ on $\mathcal{H}$. The idea is to compare two sheaves $R^2\pi^{an}_*\mathbf{C}$ and $\mathcal{H}$. Note that, for each point $p \in \Sigma$, the germ (U, p) is isomorphic to the product of an ADE surface singularity S and $(\mathbf{C}^{2n-2}, 0)$. Let $\tilde{S}$ be the minimal resolution of S. Then, $(R^2\pi^{an}_*\mathbf{C})_p$ is isomorphic to $H^2(\tilde{S}, \mathbf{C})$. A monodromy of $R^2\pi^{an}_*\mathbf{C}$ comes from a graph automorphism of the Dynkin diagram determined by the exceptional (-2)-curves on $\tilde{S}$. As is well known, S is described in terms of a simple Lie algebra $\mathfrak{g}$, and $H^2(\tilde{S}, \mathbf{C})$ is identified with the Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$; therefore, one may regard $R^2\pi^{an}_*\mathbf{C}$ as a local system of the $\mathbf{C}$-module $\mathfrak{h}$ (on Σ), whose monodromy action coincides with the natural action of a graph automorphism on $\mathfrak{h}$. On the other hand, $\mathcal{H}$ is a local system of $\mathfrak{h}/W$, where $\mathfrak{h}/W$ is the linear space obtained as the quotient of $\mathfrak{h}$ by the Weyl group W of $\mathfrak{g}$. The action of a graph automorphism on $\mathfrak{h}$ descends to an action on $\mathfrak{h}/W$, which gives a monodromy action for $\mathcal{H}$. This description of the monodromy enables us to compute $\dim H^0(\Sigma, \mathcal{H})$.

---

[1] More exactly, this means that the Poisson deformations are locally trivial as usual flat deformations of U^{an}

Proposition (4.2) together with the exact sequence above gives an upper-bound of $\dim \mathrm{PD}_U(\mathbf{C}[\epsilon])$ in terms of some topological data of X (or U). In §5, we shall prove Theorem (5.1) by using this upper-bound. The rough idea is the following. There is a natural map of functors $\mathrm{PD}_{\tilde{U}} \to \mathrm{PD}_U$ induced by the resolution map $\tilde{U} \to U$. The tangent space $\mathrm{PD}_{\tilde{U}}(\mathbf{C}[\epsilon])$ to $\mathrm{PD}_{\tilde{U}}$ is identified with $H^2(\tilde{U}^{an}, \mathbf{C})$. We have an exact sequence

$$0 \to H^2(U^{an}, \mathbf{C}) \to H^2(\tilde{U}^{an}, \mathbf{C}) \to H^0(U^{an}, R^2\pi^{an}_*\mathbf{C}) \to 0,$$

and $\dim H^0(U^{an}, R^2\pi^{an}_*\mathbf{C}) = m$. In particular, we have $\dim H^2(\tilde{U}^{an}, \mathbf{C}) = \dim H^2(U^{an}, \mathbf{C}) + m$. But this implies that $\dim \mathrm{PD}_{\tilde{U}}(\mathbf{C}[\epsilon]) \geq \dim \mathrm{PD}_U(\mathbf{C}[\epsilon])$. On the other hand, the map $\mathrm{PD}_{\tilde{U}} \to \mathrm{PD}_U$ has a finite closed fiber; or more exactly, the corresponding map $\mathrm{Spec} R_{\tilde{U}} \to \mathrm{Spec} R_U$ of prorepresentable hulls, has a finite closed fiber. Since $\mathrm{PD}_{\tilde{U}}$ is unobstructed, this implies that PD_U is unobstructed and $\dim \mathrm{PD}_{\tilde{U}}(\mathbf{C}[\epsilon]) = \dim \mathrm{PD}_U(\mathbf{C}[\epsilon])$. Finally, we obtain the unobstructedness of PD_X from that of PD_U.

Theorem (5.1) is only concerned with the formal deformations of X; but, if we impose the following condition (*), then the formal universal Poisson deformation of X has an algebraization.

(*): X has a $\mathbf{C}^*$-action with positive weights with a unique fixed point $0 \in X$. Moreover, ω is positively weighted for the action.

We shall briefly explain how this condition (*) is used in the algebraization. Let $R_X := \lim R_X/(m_X)^{n+1}$ be the prorepresentable hull of PD_X. Then the formal universal deformation $\{X_n\}$ of X defines an m_X-adic ring $A := \lim \Gamma(X_n, \mathcal{O}_{X_n})$ and let $\hat{A}$ be the completion of A along the maximal ideal of A. The rings R_X and $\hat{A}$ both have natural $\mathbf{C}^*$-actions induced from the $\mathbf{C}^*$-action on X, and there is a $\mathbf{C}^*$-equivariant map $R_X \to \hat{A}$. By taking the $\mathbf{C}^*$-subalgebras of R_X and $\hat{A}$ generated by eigen-vectors, we get a map

$$\mathbf{C}[x_1, ..., x_d] \to S$$

from a polynomial ring to a $\mathbf{C}$-algebra of finite type. We also have a Poisson structure on S over $\mathbf{C}[x_1, ..., x_d]$ by the second condition of (*). As a consequence, there is an affine space $\mathbf{A}^d$ whose completion at the origin coincides with $\mathrm{Spec}(R_X)$ in such a way that the formal universal Poisson deformation over $\mathrm{Spec}(R_X)$ is algebraized to a $\mathbf{C}^*$-equivariant map

$$\mathcal{X} \to \mathbf{A}^d.$$

Now, by using the minimal model theory due to Birkar-Cascini-Hacon-McKernan [BCHM], one can study the general fiber of $\mathcal{X} \to \mathbf{A}^d$. According to [BCHM], we can take a crepant partial resolution $\pi : Y \to X$ in such a way that Y has only $\mathbf{Q}$-factorial terminal singularities. This Y is called a $\mathbf{Q}$-*factorial terminalization* of X. In our case, Y is a symplectic variety and the $\mathbf{C}^*$-action on X uniquely extends to that on Y. Since Y has only terminal singularities, it is relatively easy to show that the Poisson deformation functor PD_Y is unobstructed. Moreover, the formal universal Poisson deformation of Y has an algebraization over an affine space $\mathbf{A}^d$:

$$\mathcal{Y} \to \mathbf{A}^d.$$

There is a $\mathbf{C}^*$-equivariant commutative diagram

$$\begin{tikzcd}
\mathcal{Y} \arrow[r] \arrow[d] & \mathcal{X} \arrow[d] \\
\mathbf{A}^d \arrow[r, "\psi"] & \mathbf{A}^d
\end{tikzcd} \tag{1}$$

By Theorem (5.5), (a): ψ is a finite surjective map, (b): $\mathcal{Y} \to \mathbf{A}^d$ is a locally trivial deformation of Y, and (c): the induced map $\mathcal{Y}_t \to \mathcal{X}_{\psi(t)}$ is an isomorphism for a general point $t \in \mathbf{A}^d$. As an application of Theorem (5.5), we have

Corollary (5.6): *Let (X, ω) be an affine symplectic variety with the property (*). Then the following are equivalent.*

(1) *X has a crepant projective resolution.*

(2) *X has a smoothing by a Poisson deformation.*

Example (i) Let $O \subset \mathfrak{g}$ be a nilpotent orbit of a complex simple Lie algebra. Let $\tilde{O}$ be the normalization of the closure $\bar{O}$ of O in $\mathfrak{g}$. Then $\tilde{O}$ is an affine symplectic variety with the Kostant-Kirillov 2-form ω on O. Let G be a complex algebraic group with $Lie(G) = \mathfrak{g}$. By [Fu], $\tilde{O}$ has a crepant projective resolution if and only if O is a Richardson orbit (cf. [C-M]) and there is a parabolic subgroup P of G such that its Springer map $T^*(G/P) \to \tilde{O}$ is birational. In this case, every crepant resolution of $\tilde{O}$ is actually obtained as a Springer map for some P. If $\tilde{O}$ has a crepant resolution, $\tilde{O}$ has a smoothing by a Poisson deformation. The smoothing of $\tilde{O}$ is isomorphic to the affine variety G/L, where L is the Levi subgroup of P. Conversely, if $\tilde{O}$ has a smoothing by a Poisson deformation, then the smoothing always has this form.

(ii) In general, $\tilde{O}$ has no crepant resolutions. But a suitable generalized Springer map gives a $\mathbf{Q}$-factorial terminalization of $\tilde{O}$ by [Na 4] and [Fu 2]. More explicitly, there is a parabolic subalgebra $\mathfrak{p}$ with Levi decomposition $\mathfrak{p} = \mathfrak{n} \oplus \mathfrak{l}$ and a nilpotent orbit O' in $\mathfrak{l}$ so that the generalized Springer map $G \times^P (\mathfrak{n} + \bar{O}') \to \tilde{O}$ is a crepant, birational map, and the normalization of $G \times^P (\mathfrak{n} + \bar{O}')$ is a $\mathbf{Q}$-factorial terminalization of $\tilde{O}$. By a Poisson deformation, $\tilde{O}$ deforms to the normalization of $G \times^L \bar{O}'$. Here $G \times^L \bar{O}'$ is a fiber bundle over G/L with a typical fiber $\bar{O}'$, and its normalization can be written as $G \times^L \tilde{O}'$ with the normalization $\tilde{O}'$ of $\bar{O}'$.

1 Local system associated with a symplectic variety

(1.1) A *symplectic variety* (X, ω) is a pair of a normal algebraic variety X defined over $\mathbf{C}$ and a symplectic 2-form ω on the regular part X_{reg} of X such that, for any resolution $\mu : \tilde{X} \to X$, the 2-form ω on $\mu^{-1}(X_{reg})$ extends to a closed regular 2-form on $\tilde{X}$. We also have a similar notion of a symplectic variety in the complex analytic category (eg. the germ of a normal complex space, a holomorphically convex, normal, complex space). For an algebraic variety X over $\mathbf{C}$, we denote by X^{an} the associated complex space. Note that if (X, ω) is a symplectic variety, then X^{an} is naturally a symplectic variety in the complex analytic category. A symplectic variety X (resp. X^{an}) has rational Gorenstein singularities. The symplectic 2-form ω defines a bivector $\Theta \in \wedge^2 \Theta_{X_{reg}}$ by the identification $\Omega^2_{X_{reg}} \cong \wedge^2 \Theta_{X_{reg}}$ by ω. Define a Poisson structure $\{\ ,\ \}$ on X_{reg} by $\{f, g\} := \Theta(df \wedge dg)$. Since X is normal, the Poisson structure on X_{reg} uniquely extends to a Poisson structure on X. Here, we recall the definition of a Poisson scheme or a Poisson complex space.

Definition. Let T be a scheme (resp. complex space). Let X be a scheme (resp. complex space) over T. Then $(X, \{\ ,\ \})$ is a Poisson scheme (resp. a Poisson space) over T if $\{\ ,\ \}$ is an $\mathcal{O}_T$-linear map:

$$\{\ ,\ \} : \wedge^2_{\mathcal{O}_T} \mathcal{O}_X \to \mathcal{O}_X$$

such that, for $a, b, c \in \mathcal{O}_X$,

1. $\{a, \{b, c\}\} + \{b, \{c, a\}\} + \{c, \{a, b\}\} = 0$
2. $\{a, bc\} = \{a, b\}c + \{a, c\}b$.

Let $(X, \{\ ,\ \})$ be a Poisson scheme (resp. Poisson space) over $\mathbf{C}$. Let S be a local Artinian $\mathbf{C}$-algebra with $S/m_S = \mathbf{C}$. Let T be the affine scheme (resp. complex space) whose coordinate ring is S. A Poisson deformation of $(X, \{\ ,\ \})$ over S is a Poisson scheme (resp. Poisson complex space) over T: $(\mathcal{X}, \{\ ,\ \}_T)$ such that $\mathcal{X}$ is flat over T, $\mathcal{X} \times_T \mathrm{Spec}(\mathbf{C}) \cong X$, and the Poisson structure $\{\ ,\ \}_T$ induces the original Poisson structure $\{\ ,\ \}$ over the closed fiber X. We define $\mathrm{PD}_X(S)$ to be the set of equivalence classes of the pairs of Poisson deformations $\mathcal{X}$ of X over $\mathrm{Spec}(S)$ and Poisson isomorphisms $\phi : \mathcal{X} \times_{\mathrm{Spec}(S)} \mathrm{Spec}(\mathbf{C}) \cong X$. Here $(\mathcal{X}, \phi)$ and $(\mathcal{X}', \phi')$ are equivalent if there is a Poisson isomorphism $\varphi : \mathcal{X} \cong \mathcal{X}'$ over $\mathrm{Spec}(S)$ which induces the identity map of X over $\mathrm{Spec}(\mathbf{C})$ via ϕ and ϕ'. We define the *Poisson deformation functor*:

$$\mathrm{PD}_{(X, \{\ ,\ \})} : (\mathrm{Art})_{\mathbf{C}} \to (\mathrm{Set})$$

from the category of local Artin $\mathbf{C}$-algebras with residue field $\mathbf{C}$ to the category of sets. Let $\mathbf{C}[\epsilon]$ be the ring of dual numbers over $\mathbf{C}$. Then the set $\mathrm{PD}_X(\mathbf{C}[\epsilon])$ has a structure of the $\mathbf{C}$-vector space, and it is called the tangent space of PD_X. A Poisson deformation of X over $\mathrm{Spec}\mathbf{C}[\epsilon]$ is particularly called a *1-st order* Poisson deformation of X. It is easy to see that $\mathrm{PD}_{(X, \{\ ,\ \})}$ satisfies the Schlessinger's conditions ([Sch]) except that possibly $\dim \mathrm{PD}_{(X, \{\ ,\ \})}(\mathbf{C}[\epsilon]) = \infty$. For details on Poisson deformations, see [G-K], [Na 2].

(1.2) Let $(S, 0)$ be the germ of a rational double point of dimension 2. More explicitly,

$$S := \{(x, y, z) \in \mathbf{C}^3; f(x, y, z) = 0\},$$

where

$$f(x, y, z) = xy + z^{r+1},$$

$$f(x, y, z) = x^2 + y^2 z + z^{r-1},$$

$$f(x, y, z) = x^2 + y^3 + z^4,$$

$$f(x, y, z) = x^2 + y^3 + yz^3,$$

or

$$f(x, y, z) = x^2 + y^3 + z^5$$

according as S is of type A_r, D_r $(r \geq 4)$ E_6, E_7 or E_8. We put

$$\omega_S := res(dx \wedge dy \wedge dz/f).$$

Then ω_S is a symplectic 2-form on $S - \{0\}$ and $(S, 0)$ becomes a symplectic variety. Let us denote by $\omega_{\mathbf{C}^{2m}}$ the canonical symplectic form on $\mathbf{C}^{2m}$:

$$ds_1 \wedge dt_1 + ... + ds_m \wedge dt_m.$$

Let (X, ω) be a symplectic variety of dimension $2n$ whose singularities are (analytically) locally isomorphic to $(S, 0) \times (\mathbf{C}^{2n-2}, 0)$. Let Σ be the singular locus of X.

Lemma (1.3) *For any $p \in \Sigma$, there are an open neighborhood $U \subset X^{an}$ of p and an open immersion*

$$\phi : U \to S \times \mathbf{C}^{2n-2}$$

such that $\omega|_U = \phi^*((p_1)^*\omega_S + (p_2)^*\omega_{\mathbf{C}^{2n-2}})$, where p_i are i-th projections of $S \times \mathbf{C}^{2n-2}$.

Proof. Let ω_1 be an arbitrary symplectic 2-form on the regular locus of $(S, 0) \times (\mathbf{C}^{2n-2}, 0)$. On the other hand, we put

$$\omega_0 := (p_1)^*\omega_S + (p_2)^*\omega_{\mathbf{C}^{2n-2}}.$$

The singularity $(S, 0)$ can be written as $(\mathbf{C}^2, 0)/G$ with a finite subgroup $G \subset SL(2, \mathbf{C})$. Let $\pi : (\mathbf{C}^2, 0) \to (S, 0)$ be the quotient map. The finite group G acts on $(\mathbf{C}^2, 0) \times (\mathbf{C}^{2n-2}, 0)$ in such a way that it acts on the second factor trivially. Then one has the quotient map

$$\pi \times id : (\mathbf{C}^2, 0) \times (\mathbf{C}^{2n-2}, 0) \to (S, 0) \times (\mathbf{C}^{2n-2}, 0).$$

We put

$$\tilde{\omega}_i := (\pi \times id)^*\omega_i$$

for $i = 0, 1$. Then $\tilde{\omega}_i$ are G-invariant symplectic 2-forms on $(\mathbf{C}^2, 0) \times (\mathbf{C}^{2n-2}, 0)$. We shall prove that there is a G-equivarinat automorphism $\tilde{\varphi}$ of $(\mathbf{C}^2, 0) \times (\mathbf{C}^{2n-2}, 0)$ such that $\tilde{\varphi}^*\tilde{\omega}_1 = \tilde{\omega}_0$. The basic idea of the following arguments is due to [Mo]. Let (x, y) be the coordinates of $(\mathbf{C}^2, 0)$ and let

$(s_1, ..., s_{n-1}, t_1, ..., t_{n-1})$ be the coordinates of $(\mathbf{C}^{2n-2}, 0)$. The symplectic 2-forms $\tilde{\omega}_0$ and $\tilde{\omega}_1$ restrict respectively to give 2-forms $\tilde{\omega}_0(\mathbf{0})$ and $\tilde{\omega}_1(\mathbf{0})$ on the tangent space $T_{\mathbf{C}^{2n},\mathbf{0}}$ at the origin $\mathbf{0} \in \mathbf{C}^{2n}$. By the definition of $\tilde{\omega}_0$,

$$\tilde{\omega}_0(\mathbf{0}) = adx \wedge dy + \Sigma ds_i \wedge dt_i$$

with some $a \in \mathbf{C}^*$. Next write $\tilde{\omega}_1(\mathbf{0})$ by using dx, dy, ds_i and dt_j. We may assume that G contains a diagonal matrix

$$\begin{pmatrix} \zeta & 0 \\ 0 & \zeta^{-1} \end{pmatrix}$$

where ζ is a primitive l-th root of unity with some $l > 1$. Since $\tilde{\omega}_1$ is G-invariant, $\tilde{\omega}_1(\mathbf{0})$ does not contain the terms $dx \wedge ds_i$, $dx \wedge dt_j$, $dy \wedge ds_i$ or $dy \wedge dt_j$. One can choose a scalar multiplication $c : (\mathbf{C}^2, 0) \to (\mathbf{C}^2, 0)$ $((x, y) \to (cx, cy))$ and a linear automorphism $\sigma : (\mathbf{C}^{2n-2}, 0) \to (\mathbf{C}^{2n-2}, 0)$ so that $\tilde{\omega}_2 := (c \times \sigma)^*(\tilde{\omega}_1)$ satisfies

$$\tilde{\omega}_2(\mathbf{0}) = adx \wedge dy + \Sigma ds_i \wedge dt_i.$$

Note that

$$\tilde{\omega}_0(\mathbf{0}) = \tilde{\omega}_2(\mathbf{0}).$$

Since $c \times \sigma$ is G-equivariant, $\tilde{\omega}_2$ is a G-invariant symplectic 2-form. For $\tau \in \mathbf{R}$, define

$$\omega(\tau) := (1 - \tau)\tilde{\omega}_0 + \tau\tilde{\omega}_2.$$

We put

$$u := d\omega(\tau)/d\tau.$$

Since $S \times \mathbf{C}^{2n-2}$ has only quotient singularities, the complex $((\pi \times id)^G_* \Omega^{\cdot}_{\mathbf{C}^2 \times \mathbf{C}^{2n-2}}, d)$ is a resolution of the constant sheaf $\mathbf{C}$ on $S \times \mathbf{C}^{2n-2}$. Note that u is a section of $(\pi \times id)^G_* \Omega^2_{\mathbf{C}^2 \times \mathbf{C}^{2n-2}}$. Moreover, u is d-closed. Therefore, one can write $u = dv$ with a G-invariant 1-form v. Moreover v can be chosen such that $v(\mathbf{0}) = 0$. Define a vector field X_τ on $(\mathbf{C}^{2n}, 0)$ by

$$i_{X_\tau}\omega(\tau) = -v.$$

Since $\omega(\tau)$ is d-closed, we have

$$L_{X_\tau}\omega(\tau) = -u$$

where $L_{X_\tau}\omega(\tau)$ is the Lie derivative of $\omega(\tau)$ along X_τ. If we take a sufficiently small open subset V of $\mathbf{0} \in \mathbf{C}^{2n}$, then the vector fields $\{X_\tau\}_{0 \le \tau \le 1}$ define a family of open immersions $\varphi_\tau : V \to \mathbf{C}^{2n}$ via

$$d\varphi_\tau / d\tau = X_\tau(\varphi_\tau), \quad \varphi_0 = id.$$

Since all φ_τ fix the origin and X_τ are all G-invariant, φ_τ induce G-equivariant automorphisms of $(\mathbf{C}^{2n}, 0)$. By the definition of X_τ, we have $(\varphi_\tau)^*\omega(\tau) = \omega(0)$. In particular, $(\varphi_1)^*\tilde{\omega}_2 = \tilde{\omega}_0$. We put

$$\tilde{\varphi} := (\varphi_1) \circ (c \times \sigma).$$

The G-equivariant automorphism $\tilde{\varphi}$ of $(\mathbf{C}^{2n}, 0)$ descends to an automorphism φ of $(S, 0) \times (\mathbf{C}^{2n-2}, 0)$ so that $\varphi^*\omega_1 = \omega_0$. Q.E.D.

We cover the singular locus Σ by a family of open sets $\{U_\alpha\}$ of X^{an} in such a way that each U_α admits an open immersion ϕ_α as in Lemma (1.3). In the remainder, we call such a covering $\{U_\alpha\}$ *admissible.*

(1.4) Let (X, ω) be the same as above. Denote by $T^1_{X^{an}}$ the analytic coherent sheaf $\underline{\mathrm{Ext}}^1(\Omega^1_{X^{an}}, \mathcal{O}_{X^{an}})$. Note that the sheaf $T^1_{X^{an}}$ is the sheafication of the presheaf associating to each open set $V \subset X^{an}$ the $\mathbf{C}$-vector space of the isomorphic classes of 1-st order deformations of V. Let us consider the presheaf on X^{an} which associates to each open set V the $\mathbf{C}$-vector space of the isomorphic classes of 1-st order *Poisson* deformation. Denote by $PT^1_{X^{an}}$ the sheafication of this presheaf. Note that both sheaves $T^1_{X^{an}}$ and $PT^1_{X^{an}}$ have support on Σ. One has a natural map

$$PT^1_{X^{an}} \to T^1_{X^{an}}$$

of sheaves of $\mathbf{C}$-modules by forgetting the Poisson structure. Define a subsheaf $\mathcal{H}$ of $T^1_{X^{an}}$ as the image of this map.

Lemma (1.5) *$\mathcal{H}$ is a locally constant $\mathbf{C}$-module over Σ.*

Proof. Take an admissible covering $\{U_\alpha\}$. For each α,

$$T^1_{U_\alpha} = (p_1 \circ \phi_\alpha)^* T^1_S.$$

We put

$$H_\alpha := (p_1 \circ \phi_\alpha)^{-1} T^1_S.$$

Note that H_α is a constant $\mathbf{C}$-module on $U_\alpha \cap \Sigma$. We shall prove that $\mathcal{H}|_{U_\alpha} = H_\alpha$. In fact, let $\mathcal{U}_\alpha \to \mathrm{Spec}\mathbf{C}[\epsilon]$ be a 1-st order Poisson deformation of U_α.

Let $0 \in U_\alpha$ be the point which corresponds to $(0,0) \in S \times \mathbf{C}^{2n-2}$ via ϕ_α. By applying the second statement of the next Lemma (1.6) to $\hat{\mathcal{O}}_{U_\alpha,0}$ and $\hat{\mathcal{O}}_{\mathcal{U}_\alpha,0}$, we conclude that $(\mathcal{U}_\alpha, 0) \cong (\mathcal{S}, 0) \times (\mathbf{C}^{2n-2}, 0)$, where $\mathcal{S}$ is a 1-st order deformation of S. Conversely, a 1-st order deformation of this form always comes from a Poisson deformation of U_α. Q.E.D.

Lemma (1.6). *Let $S := \{f(x,y,z) = 0\} \subset \mathbf{C}^3$ be an isolated hypersurface singularity which admits a Poisson structure, and let $(\mathbf{C}^{2n-2}, 0)$ be a symplectic manifold with the standard symplectic structure. Put $V := (S,0) \times (\mathbf{C}^{2n-2}, 0)$ and introduce the product Poisson structure on V. Assume that $\mathcal{V} \to \mathrm{Spec}\, \mathbf{C}[\epsilon]$ is a 1-st order Poisson deformation of V. Then*

$$\mathcal{V} \cong (\mathcal{S}, 0) \times (\mathbf{C}^{2n-2}, 0)$$

as a flat deformation. Here $(\mathcal{S}, 0)$ is a 1-st order flat deformation of $(S, 0)$.

Proof. We denote by $\mathbf{s} = (s_1, ..., s_{2n-2})$ the coordinates of $\mathbf{C}^{2n-2}$. Let $f_1, ..., f_\tau \in \mathbf{C}\{x,y,z\}$ be the representatives of a basis of $\mathbf{C}\{x,y,z\}/(f, f_x, f_y, f_z)$. The 1-st order deformation $\mathcal{V}$ can be written as

$$f(x,y,z) + \epsilon(f_1(x,y,z)g_1(\mathbf{s}) + ... + f_\tau(x,y,z)g_\tau(\mathbf{s})) = 0.$$

We prove that g_i are all constants. Let $\{\ ,\ \}$ be the Poisson structure on V. By the definition, we have

$$\{x, s_i\} = \{y, s_i\} = \{z, s_i\} = 0$$

in $\mathcal{O}_{V,0}$. Let $\{\ ,\ \}'$ be the Poisson structure on $\mathcal{V}$ extending the Poisson structure $\{\ ,\ \}$. Then we have

$$\{x, s_i\}' = \epsilon\alpha_i,\ \{y, s_i\}' = \epsilon\beta_i,\ \{z, s_i\}' = \epsilon\gamma_i,$$

for some elements α_i, β_i and γ_i in $\mathcal{O}_{V,0}$. Since $f + \epsilon(f_1g_1 + ... + f_\tau g_\tau) = 0$ in $\mathcal{O}_{\mathcal{V},0}$, we must have

$$\{f + \epsilon(f_1g_1 + ... + f_\tau g_\tau), s_i\}' = 0$$

in $\mathcal{O}_{\mathcal{V},0}$. By calculating the left-hand side, one has

$$f_x\{x, s_i\}' + f_y\{y, s_i\}' + f_z\{z, s_i\}' + \epsilon(\Sigma_{1\leq j\leq\tau} f_j\{g_j, s_i\} + \Sigma_{1\leq j\leq\tau} g_j\{f_j, s_i\}) = 0$$

Recall that

$$\{s_1, s_2\} = \{s_3, s_4\} = ... = \{s_{2n-3}, s_{2n-2}\} = 1,$$

and $\{s_k, s_l\} = 0$ for other $k < l$. Moreover, note that $\{f_j, s_i\} = 0$. Assume that i is odd, then one has

$$\epsilon(f_x\alpha_i + f_y\beta_i + f_z\gamma_i + \Sigma_{1\leq i\leq\tau} f_j \cdot (g_j)_{s_{i+1}}) = 0.$$

This implies that

$$f_x\alpha_i + f_y\beta_i + f_z\gamma_i + \Sigma_{1\leq i\leq\tau} f_j \cdot (g_j)_{s_{i+1}} = 0$$

in $\mathcal{O}_{V,0}$. Note that $\mathcal{O}_{V,0} = \mathbf{C}\{x, y, z, \mathbf{s}\}/(f)$. Let us consider the equation in $\mathbf{C}\{x, y, z, \mathbf{s}\}/(f, f_x, f_y, f_z)$. Then we have

$$\Sigma_{1\leq j\leq\tau} f_j \cdot (g_j)_{s_{i+1}} = 0.$$

This implies that $(g_j)_{s_{i+1}} = 0$ in $\mathbf{C}\{\mathbf{s}\}$ for all j. When i is even, a similar argument shows that $(g_j)_{s_{i-1}} = 0$ for all j. As a consequence, g_j are constants for all j. Q.E.D.

(1.7) **Monodromy of $\mathcal{H}$**

Let γ be a closed loop in Σ starting from $p \in \Sigma$. We shall describe the monodromy of $\mathcal{H}$ along γ in terms of a certain symplectic automorphism of the germ (X^{an}, p). In order to do this, we take a sequence of admissible open sets of X^{an}: U_1, ..., U_k, $U_{k+1} := U_1$ in such a way that $p \in U_1$, $\gamma \subset \cup U_i$, $U_i \cap U_{i+1} \cap \gamma \neq \emptyset$ for $i = 1, ..., k$. Put $p_1 := p$ and choose a point $p_i \in U_i \cap U_{i+1} \cap \gamma$ for each $i \geq 2$. Let $\phi_i : U_i \to S \times \mathbf{C}^{2n-2}$ be the symplectic open immersion associated with the admissible open subset U_i. Since $\mathcal{H}$ is a locally constant $\mathbf{C}$-module by (1.5), an element of $\mathcal{H}_{p_i}$ uniquely extends to a section of $\mathcal{H}$ over U_i. Since $p_{i-1} \in U_i$, this section restricts to give an element of $\mathcal{H}_{p_{i-1}}$. In this way, we have an identification

$$m_i : \mathcal{H}_{p_{i-1}} \cong \mathcal{H}_{p_i}$$

for each i. The monodromy transformation m_γ is the composite of m_i's:

$$m_\gamma = m_{k+1} \circ ... \circ m_2.$$

One can describe each m_i in terms of certain symplectic isomorphisms as explained below. Since U_i contains p_i, the germ (X^{an}, p_i) is identified with $(S \times \mathbf{C}^{2n-2}, \phi_i(p_i))$ by ϕ_i. On the other hand, since U_i contains p_{i-1}, the germ (X^{an}, p_{i-1}) is identified with $(S \times \mathbf{C}^{2n-2}, \phi_i(p_{i-1}))$. Note that $\phi_i(p_i) = (0, *) \in S \times \mathbf{C}^{2n-2}$ and $\phi_i(p_{i-1}) = (0, **) \in S \times \mathbf{C}^{2n-2}$ for some points $*, ** \in \mathbf{C}^{2n-2}$

because $p_i, p_{i-1} \in \gamma$. Denote by $\sigma_i : \mathbf{C}^{2n-2} \to \mathbf{C}^{2n-2}$ the translation map such that $\sigma_i(*) = **$. Then, by the automorphism $id \times \sigma_i$ of $S \times \mathbf{C}^{2n-2}$, two germs $(S \times \mathbf{C}^{2n-2}, \phi_i(p_i))$ and $(S \times \mathbf{C}^{2n-2}, \phi_i(p_{i-1}))$ are identified. As a consequence, two germs (X^{an}, p_{i-1}) and (X^{an}, p_i) have been identified. By definition, this identification preserves the natural symplectic forms on (X^{an}, p_{i-1}) and (X^{an}, p_i). The symplectic isomorphism $(X^{an}, p_{i-1}) \cong (X^{an}, p_i)$ determines an isomorphism $\mathcal{H}_{p_{i-1}} \cong \mathcal{H}_{p_i}$. It is easy to see that this isomorphism coincides with m_i defined above. Note that the symplectic automorphism depends on the choice of ϕ_i, but m_i is independent of it. Now the sequence of identifications $(X^{an}, p_1) \cong (X^{an}, p_2)$, $(X^{an}, p_2) \cong (X^{an}, p_3)$, ..., $(X^{an}, p_k) \cong (X^{an}, p_1)$ finally defines a symplectic automorphism

$$i_\gamma : (X^{an}, p) \cong (X^{an}, p).$$

The map i_γ induces an automorphism of $\mathcal{H}_p$, which coincides with m_γ because $m_\gamma = m_{k+1} \circ ... \circ m_2$. Although i_γ depends on the choices of ϕ_i's, m_γ is independent of them by the definition.

(1.8) In the above, we only considered a symplectic variety whose singularities are locally isomorphic to $(S, 0) \times (\mathbf{C}^{2n-2}, 0)$. From now on, we will treat a general symplectic variety (X, ω). Let $U \subset X$ be the locus where X is smooth, or is locally a trivial deformation of a (surface) rational double point. Put $\Sigma := \mathrm{Sing}(U)$. As an open set of X, U naturally becomes a Poisson scheme. Since $X \setminus U$ has codimension at least 4 in X ([Ka 1]), one can prove in the same way as [Na 2, Proposition 13] that

$$\mathrm{PD}_X(\mathbf{C}[\epsilon]) \cong \mathrm{PD}_U(\mathbf{C}[\epsilon]).$$

Let $\mathrm{PD}_{lt,U}$ be the locally trivial Poisson deformation functor of U. More exactly, $\mathrm{PD}_{lt,U}$ is the subfunctor of PD_U corresponding to the Poisson deformations of U which are locally trivial as flat deformations of U^{an} (after forgetting Poisson structure). We shall insert a lemma here, which will be used in the proof of Proposition (1.11).

Lemma (1.9) *Let X be an affine symplectic variety let $j : X_{reg} \to X$ be the open immersion of the regular part X_{reg} into X. Then*

$$\mathrm{PD}_{lt,X}(\mathbf{C}[\epsilon]) = \mathbf{H}^2(\Gamma(X, j_*(\wedge^{\geq 1}\Theta_{X^{reg}})),$$

where $(\wedge^{\geq 1}\Theta_{X^{reg}}, \delta)$ is the Lichnerowicz-Poisson complex for X_{reg} (cf. [Na 2, §2]).

Proof. The 2-nd cohomology $\mathbf{H}^2(\Gamma(X_{reg}, \wedge^{\geq 1}\Theta_{X_{reg}}))$ describes the equivalence classes of the extension of the Poisson structure $\{\ ,\ \}$ on X_{reg} to that on $X_{reg} \times \mathrm{Spec}\mathbf{C}[\epsilon] \to \mathrm{Spec}\mathbf{C}[\epsilon]$. In fact, for $\psi \in \Gamma(X_{reg}, \wedge^2\Theta_{X_{reg}})$, we define a Poisson structure $\{\ ,\ \}_\epsilon$ on $\mathcal{O}_{X_{reg}} \oplus \epsilon\mathcal{O}_{X_{reg}}$ by

$$\{f + \epsilon f', g + \epsilon g'\}_\epsilon := \{f, g\} + \epsilon(\psi(df \wedge dg) + \{f, g'\} + \{f', g\}).$$

Then this bracket is a Poisson bracket if and only if $\delta(\psi) = 0$. On the other hand, an element $\theta \in \Gamma(X_{reg}, \Theta_{X_{reg}})$ corresponds to an automorphism φ_θ of $X_{reg} \times \mathrm{Spec}\mathbf{C}[\epsilon]$ over $\mathrm{Spec}\mathbf{C}[\epsilon]$ which restricts to give the identity map of the closed fiber X_{reg}. Let $\{\ ,\ \}_\epsilon$ and $\{\ ,\ \}'_\epsilon$ be the Poisson structures determined respectively by $\psi \in \Gamma(X_{reg}, \wedge^2\Theta_{X_{reg}})$ and $\psi' \in \Gamma(X_{reg}, \wedge^2\Theta_{X_{reg}})$. Then the two Poisson structures are equivalent under φ_θ if and only if $\psi - \psi' = \delta(\theta)$. For an affine variety X, a locally trivial infinitesimal deformation is nothing but a trivial infinitesimal deformation because $H^1(X, \Theta_X) = 0$. The original Poisson structure on X restricts to give a Poisson structure on X_{reg}. As seen above, its extension to $X_{reg} \times \mathrm{Spec}\mathbf{C}[\epsilon]$ is classified by $\mathbf{H}^2(\Gamma(X_{reg}, \wedge^{\geq 1}\Theta_{X_{reg}}))$. Each Poisson structure on $X_{reg} \times \mathrm{Spec}\mathbf{C}[\epsilon]$ can extend uniquely to that on $X \times \mathrm{Spec}\mathbf{C}[\epsilon]$.

Remark (1.10). By the same argument as [Na 2], Proposition 8, one can prove that, for a (non-affine) symplectic variety X,

$$\mathrm{PD}_{lt,X}(\mathbf{C}[\epsilon]) = \mathbf{H}^2(X, j_*(\wedge^{\geq 1}\Theta_{X_{reg}})),$$

where $\mathbf{H}^2$ is the 2-nd hypercohomology.

Let us return to the original situation in (1.8). Let $\mathcal{H} \subset T^1_{U^{an}}$ be the local constant $\mathbf{C}$-modules over Σ. We have an exact sequence of $\mathbf{C}$-vector spaces:

$$0 \to \mathrm{PD}_{lt,U}(\mathbf{C}[\epsilon]) \to \mathrm{PD}_U(\mathbf{C}[\epsilon]) \to H^0(\Sigma, \mathcal{H}).$$

The following proposition shows that the tangent space of the Poisson deformation functor of an affine symplectic variety is finite dimensional.

Proposition (1.11). *Assume that X is an affine symplectic variety. Then*

$$\mathrm{PD}_{lt,U}(\mathbf{C}[\epsilon]) \cong H^2(U^{an}, \mathbf{C}).$$

In particular, $\dim \mathrm{PD}_X(\mathbf{C}[\epsilon]) < \infty$.

Proof. Let U^0 be the smooth part of U and let $j : U^0 \to U$ be the inclusion map. Let $(\wedge^{\geq 1}\Theta_{U^0}, \delta)$ be the Lichnerowicz-Poisson complex for U^0. By Remark (1.10), one has

$$\mathrm{PD}_{lt,U}(\mathbf{C}[\epsilon]) \cong \mathbf{H}^2(U, j_*(\wedge^{\geq 1}\Theta_{U_0})).$$

By the symplectic form ω, the complex $(j_*(\wedge^{\geq 1}\Theta_{U_0}), \delta)$ is identified with $\{j_*(\wedge^{\geq 1}\Omega^1_{U_0}), d)$ (cf. [Na 2, Proposition 9]). The latter complex is the truncated *de Rham complex for a V-manifold U* $(\tilde{\Omega}^{\geq 1}_U, d)$ (cf. [St]). Let us consider the distinguished triangle

$$\tilde{\Omega}^{\geq 1}_U \to \tilde{\Omega}^{\cdot}_U \to \mathcal{O}_U \to \tilde{\Omega}^{\geq 1}_U[1].$$

We have an exact sequence

$$H^1(\mathcal{O}_U) \to \mathbf{H}^2(\tilde{\Omega}^{\geq 1}_U) \to \mathbf{H}^2(\tilde{\Omega}^{\cdot}_U) \to H^2(\mathcal{O}_U).$$

Since X is a symplectic variety, X is Cohen-Macaulay (cf. (1.1)). Moreover, X is affine and $X \setminus U$ has codimension ≥ 4 in X. Thus, by a depth argument, we see that $H^1(\mathcal{O}_U) = H^2(\mathcal{O}_U) = 0$. On the other hand, by Grothendieck's theorem [Gr][2] for V-manifolds, we have $\mathbf{H}^2(\tilde{\Omega}^{\cdot}_U) \cong \mathbf{H}^2(U^{an}, \mathbf{C})$. Now the result follows from the exact sequence above. Q.E.D.

2 Prorepresentability of the Poisson deformation functors

Let $(X, \{\ ,\ \})$ be a Poisson scheme. In this section, we shall prove that, in many important cases, $\mathrm{PD}_{(X,\{\ ,\ \})}$ has a prorepresentable hull R_X (cf. [Sch]),

---

[2] The V-manifold case is reduced to the smooth case as follows. Let W be an algebraic variety with quotient singularities (V-manifold). One can cover W by finite affine open subsets U_i, $0 \leq i \leq n$ so that each U_i admits an etale Galois cover U'_i such that $U'_i = V_i/G_i$ with a smooth variety V_i and a finite group G_i. It can be checked that, for each intersection $U_{i_0,...,i_p} := U_{i_0} \cap ... \cap U_{i_p}$, Grothendieck's theorem holds. Now one has Grothendieck's theorem for W by comparing two spectral sequences

$$E_1^{p,q} := \oplus_{i_0<...<i_p} H^q(U_{i_0,...,i_p}, \tilde{\Omega}^{\cdot}_{U_{i_0,...,i_p}}) \Longrightarrow H^{p+q}(W, \tilde{\Omega}^{\cdot}_W)$$

and

$$E'^{p,q}_1 := \oplus_{i_0<...<i_p} H^q(U^{an}_{i_0,...,i_p}, \mathbf{C}) \Longrightarrow H^{p+q}(W^{an}, \mathbf{C}).$$

and it is actually prorepresentable, i.e. $\mathrm{Hom}(R_X, \cdot) \cong \mathrm{PD}_{(X,\{\ ,\ \})}(\cdot)$. Let $\mathcal{X}$ be a Poisson scheme over a local Artinian base T and let X be the central closed fiber. Let $G_{\mathcal{X}/T}$ be the sheaf of automorphisms of $\mathcal{X}/T$. More exactly, it is a sheaf on X which associates to each open set $U \subset X$, the set of the automorphisms of the usual scheme $\mathcal{X}|_U$ over T which induce the identity map on the central fiber $U = X|_U$. Moreover, let $PG_{\mathcal{X}/T}$ be the sheaf of *Poisson automorphisms* of $\mathcal{X}/T$ as a subsheaf of $G_{\mathcal{X}/T}$. In order to show that $\mathrm{PD}_{(X,\{\ ,\ \})}$ is prorepresentable, it is enough to prove that $H^0(X, PG_{\mathcal{X}/T}) \to H^0(X, PG_{\bar{\mathcal{X}}/\bar{T}})$ is surjective for any closed subscheme $\bar{T} \subset T$ and $\bar{\mathcal{X}} := \mathcal{X} \times_T \bar{T}$. Assume that $\mathcal{X}$ is smooth over T. We denote by $\Theta_{\mathcal{X}/T}$ the relative tangent sheaf for $\mathcal{X} \to T$. Consider the Lichnerowicz-Poisson complex (cf. [Na 2, Section 2])

$$0 \to \Theta_{\mathcal{X}/T} \xrightarrow{\delta_1} \wedge^2\Theta_{\mathcal{X}/T} \xrightarrow{\delta_2} \wedge^3\Theta_{\mathcal{X}/T} ...$$

and define $P\Theta_{\mathcal{X}/T} := \mathrm{Ker}(\delta_1)$. We denote by $\Theta^0_{\mathcal{X}/T}$ (resp. $P\Theta^0_{\mathcal{X}/T}$) the subsheaf of $\Theta_{\mathcal{X}/T}$ (resp. $P\Theta^0_{\mathcal{X}/T}$) which consists of the sections vanishing on the central closed fiber.

Proposition (2.1)(Wavrik): *There is an isomorphism of sheaves of sets*

$$\alpha : \Theta^0_{\mathcal{X}/T} \cong G_{\mathcal{X}/T}.$$

Moreover, α *induces an injection*

$$P\Theta^0_{\mathcal{X}/T} \to PG_{\mathcal{X}/T}.$$

Proof. Each local section φ of $\Theta^0_{\mathcal{X}/T}$ is regarded as a derivation of $\mathcal{O}_{\mathcal{X}}$. Then we put

$$\alpha(\varphi) := id + \varphi + 1/2!(\varphi \circ \varphi) + 1/3!(\varphi \circ \varphi \circ \varphi) + ...$$

By using the property

$$\varphi(fg) = f\varphi(g) + \varphi(f)g,$$

one can check that $\alpha(\varphi)$ is an automorphism of $\mathcal{X}/T$ inducing the identity map on the central fiber. If φ is a local section of $P\Theta^0_{\mathcal{X}/T}$, then φ satisfies

$$\varphi(\{f, g\}) = \{f, \varphi(g)\} + \{\varphi(f), g\}.$$

By this property, one sees that $\alpha(\varphi)$ becomes a Poisson automorphism of $\mathcal{X}/T$. For the bijectivity of α, see [Wav].

Proposition (2.2). *In Proposition (2.1), if $\mathcal{X}$ is a Poisson deformation of a smooth symplectic variety (X, ω), then α induces an isomorphism*

$$P\Theta^0_{\mathcal{X}/T} \cong PG_{\mathcal{X}/T}.$$

Proof. We only have to prove that the map is surjective. We may assume that X is affine. Let S be the Artinian local ring with $T = \mathrm{Spec}(S)$ and let m be the maximal ideal of S. Put $T_n := \mathrm{Spec}(S/m^{n+1})$. The sequence

$$T_0 \subset T_1 \subset ... \subset T_k$$

terminates at some k and $T_k = T$. We put $X_n := \mathcal{X} \times_T T_n$. Let ϕ be a section of $PG_{\mathcal{X}/T}$. One can write

$$\phi|_{X_1} = id + \varphi_1$$

with $\varphi_1 \in m \cdot P\Theta_X$. By the next lemma, φ_1 lifts to some $\tilde{\varphi}_1 \in P\Theta_{\mathcal{X}/T}$. Then one can write

$$\phi|_{X_2} = \alpha(\tilde{\varphi}_1)|_{X_2} + \varphi_2$$

with $\varphi_2 \in m^2 \cdot P\Theta_X$. Again, by the lemma, φ_2 lifts to some $\tilde{\varphi}_2 \in P\Theta_{\mathcal{X}/T}$. Continue this operation and we finally conclude that

$$\phi = \alpha(\tilde{\varphi}_1 + \tilde{\varphi}_2 + ...).$$

Lemma (2.3). *Let $\mathcal{X} \to T$ be a Poisson deformation of a smooth symplectic variety (X, ω) over a local Artinian base $T = \mathrm{Spec}(S)$. Let $\bar{T} \subset T$ be a closed subscheme and put $\bar{\mathcal{X}} := \mathcal{X} \times_T \bar{T}$. Then the restriction map*

$$P\Theta_{\mathcal{X}/T} \to P\Theta_{\bar{\mathcal{X}}/\bar{T}}$$

is surjective.

Proof. We may assume that X is affine. The Lichnerowicz-Poisson complex $(\wedge^{\geq 1}\Theta_{\mathcal{X}/T}, \delta)$ is identified with the truncated de Rham complex $(\Omega^{\geq 1}_{\mathcal{X}/T}, d)$ by the symplectic 2-form ω (cf. [Na 2], Section 2). There is a distinguished triangle

$$\Omega^{\geq 1}_{\mathcal{X}/T} \to \Omega^{\cdot}_{\mathcal{X}/T} \to \mathcal{O}_{\mathcal{X}} \to \Omega^{\geq 1}_{\mathcal{X}/T}[1],$$

and it induces an exact sequence

$$... \to HP^i(\mathcal{X}/T) \to H^i(X^{an}, S) \to H^i(X, \mathcal{O}_{\mathcal{X}}) \to ...$$

In particular, we have an exaxt sequence

$$0 \to K \to HP^1(\mathcal{X}/T) \to H^1(X^{an}, S) \to 0,$$

where

$$K := \mathrm{Coker}[H^0(X^{an}, S) \to H^0(X, \mathcal{O}_{\mathcal{X}})].$$

Similarly for $\bar{\mathcal{X}}$, we have an exact sequence

$$0 \to \bar{K} \to HP^1(\bar{\mathcal{X}}/\bar{T}) \to H^1(X^{an}, \bar{S}) \to 0$$

with

$$\bar{K} := \mathrm{Coker}[H^0(X^{an}, \bar{S}) \to H^0(X, \mathcal{O}_{\bar{\mathcal{X}}})].$$

Since the restriction maps $K \to \bar{K}$ and $H^0(X^{an}, S) \to H^0(X^{an}, \bar{S})$ are both surjective, the restriction map $HP^1(\mathcal{X}/T) \to HP^1(\bar{\mathcal{X}}/\bar{T})$ is surjective. Finally, note that $HP^1(\mathcal{X}/T) = H^0(X, P\Theta_{\mathcal{X}/T})$ and $HP^1(\bar{\mathcal{X}}/\bar{T}) = H^0(X, P\Theta_{\bar{\mathcal{X}}/\bar{T}})$.

Proposition (2.4). *In the same assumption in Lemma (2.3), if the restriction map*

$$H^0(X, P\Theta_{\mathcal{X}/T}) \to H^0(X, P\Theta_{\bar{\mathcal{X}}/\bar{T}})$$

is surjective, then the restriction map

$$H^0(X, PG_{\mathcal{X}/T}) \to H^0(X, PG_{\bar{\mathcal{X}}/\bar{T}})$$

is surjective.

Proof. If the map

$$H^0(X, P\Theta_{\mathcal{X}/T}) \to H^0(X, P\Theta_{\bar{\mathcal{X}}/\bar{T}})$$

is surjective,

$$H^0(X, P\Theta^0_{\mathcal{X}/T}) \to H^0(X, P\Theta^0_{\bar{\mathcal{X}}/\bar{T}})$$

is surjective. Then the result follows from Proposition (2.2).

Corollary (2.5). *The Poisson deformation functor $PD_{(X,\{\ ,\ \})}$ for a symplectic variety (X, ω), is prorepresentable in the following two cases:*

(1) *X is convex (i.e. X has a birational projective morphism to an affine variety), and admits only terminal singularities.*

(2) *X is affine, and $H^1(X^{an}, \mathbf{C}) = 0$.*

Proof. First, we must show that $\dim \mathrm{PD}_{(X,\{\ ,\ \})}(\mathbf{C}[\epsilon]) < \infty$. Let U be the smooth part of X. In the case (1), we have $\mathrm{PD}_{(X,\{\ ,\ \})}(\mathbf{C}[\epsilon]) = H^2(U^{an}, \mathbf{C})$; hence $\mathrm{PD}_{(X,\{\ ,\ \})}(\mathbf{C}[\epsilon])$ is a finite dimensional $\mathbf{C}$-vector space. For the case (2), the finiteness is proved in Proposition (1.10). Assume that $\mathcal{X} \to T$ is a Poisson deformation of X with a local Artinian base. Let $\bar{T}$ be a closed subscheme of T and let $\bar{\mathcal{X}} \to \bar{T}$ be the induced Poisson deformation of X over $\bar{T}$. Let $\mathcal{U} \subset \mathcal{X}$ (resp. $\bar{\mathcal{U}} \subset \bar{\mathcal{X}}$) be the open locus where the map $\mathcal{X} \to T$ (resp. $\bar{\mathcal{X}} \to \bar{T}$) is smooth. Let j be the inclusion map of $\mathcal{U}$ to $\mathcal{X}$. Since $j_*\mathcal{O}_{\mathcal{U}} = \mathcal{O}_{\mathcal{X}}$, a Poisson automorphism of $\mathcal{U}$ (which induces the identity on the closed fiber) uniquely extends to that of $\mathcal{X}$. Therefore, we have an isomorphism

$$H^0(\mathcal{X}, PG_{\mathcal{X}/T}) \cong H^0(\mathcal{U}, PG_{\mathcal{U}/T}).$$

Similarly, we have

$$H^0(\bar{\mathcal{X}}, PG_{\bar{\mathcal{X}}/\bar{T}}) \cong H^0(\bar{\mathcal{U}}, PG_{\bar{\mathcal{U}}/\bar{T}}).$$

By Proposition (2.4), it suffices to show that the restriction map

$$H^0(U, P\Theta_{\mathcal{U}/T}) \to H^0(U, P\Theta_{\bar{\mathcal{U}}/\bar{T}})$$

is surjective.

For the case (1), we have already proved the surjectivity in [Na 2], Theorem 14. Let us consider the case (2). Note that $H^0(U, P\Theta_{\mathcal{U}/T}) \cong \mathbf{H}^1(U, \Theta^{\geq 1}_{\mathcal{U}/T})$, where $(\Theta^{\geq 1}_{\mathcal{U}/T}, \delta)$ is the Lichnerowicz-Poisson complex for $\mathcal{U}/T$. As in the proof of Lemma (2.3), the Lichnerowicz-Poisson complex is identified with the truncated de Rham complex $(\Omega^{\geq 1}_{\mathcal{U}/T}, d)$, and it induces the exact sequence

$$0 \to K \to \mathbf{H}^1(U, \Omega^{\geq 1}_{\mathcal{U}/T}) \to H^1(U^{an}, S),$$

where S is the affine ring of T, and $K := \mathrm{Coker}[H^0(U^{an}, S) \to H^0(U, \mathcal{O}_{\mathcal{U}})]$. We shall prove that $H^1(U^{an}, S) = 0$. Since $H^1(U^{an}, S) = H^1(U^{an}, \mathbf{C}) \otimes S$, it suffices to show that $H^1(U^{an}, \mathbf{C}) = 0$. Let $f : \tilde{X} \to X$ be a resolution of X such that $f^{-1}(U) \cong U$ and the exceptional locus E of f is a divisor with only simple normal crossing. One has the exact sequence

$$H^1(\tilde{X}^{an}, \mathbf{C}) \to H^1(U^{an}, \mathbf{C}) \to H^2_E(\tilde{X}^{an}, \mathbf{C}) \to H^2(\tilde{X}^{an}, \mathbf{C}),$$

where the first term is zero because X has only rational singularities and $H^1(X^{an}, \mathbf{C}) = 0$. We have to prove that $H^2_E(\tilde{X}^{an}, \mathbf{C}) \to H^2(\tilde{X}^{an}, \mathbf{C})$ is an injection. Put $n := \dim X$; then, $H^2_E(\tilde{X}^{an}, \mathbf{C})$ is dual to the cohomology $H^{2n-2}_c(E^{an}, \mathbf{C})$ with compact support (cf. the proof of Proposition 2 of [Na 3]). Let $E = \cup E_i$ be the irreducible decomposition of E. The p-multiple locus of E is, by definition, the locus of points of E which are contained in the intersection of some p different irreducible components of E. Let $E^{[p]}$ be the normalization of the p-multiple locus of E. For example, $E^{[1]}$ is the disjoint union of E_i's, and $E^{[2]}$ is the normalization of the singular locus of E. There is an exact sequence

$$0 \to \mathbf{C}_E \to \mathbf{C}_{E^{[1]}} \to \mathbf{C}_{E^{[2]}} \to ...$$

By using this exact sequence, we see that $H^{2n-2}_c(E^{an}, \mathbf{C})$ is a $\mathbf{C}$-vector space whose dimension equals the number of irreducble components of E. By the duality, we have

$$H^2_E(\tilde{X}^{an}, \mathbf{C}) = \oplus \mathbf{C}[E_i]$$

and the map $H^2_E(\tilde{X}^{an}, \mathbf{C}) \to H^2(\tilde{X}^{an}, \mathbf{C})$ is an injection. Therefore, $H^1(U^{an}, \mathbf{C}) = 0$. We now know that

$$H^0(U, P\Theta_{\mathcal{U}/T}) \cong K.$$

Similarly, we have

$$H^0(U, P\Theta_{\bar{\mathcal{U}}/\bar{T}}) \cong \bar{K},$$

where $\bar{K} := \mathrm{Coker}[H^0(U, \bar{S}) \to H^0(U, \mathcal{O}_{\bar{\mathcal{U}}})]$ and $\bar{S}$ is the affine ring of $\bar{T}$. Since the restriction maps $H^0(X, \mathcal{O}_{\mathcal{X}}) \to H^0(U, \mathcal{O}_{\mathcal{U}})$ and $H^0(X, \mathcal{O}_{\bar{\mathcal{X}}}) \to H^0(U, \mathcal{O}_{\bar{\mathcal{U}}})$ are both isomorphisms, the restriction map $H^0(U, \mathcal{O}_{\mathcal{U}}) \to H^0(U, \mathcal{O}_{\bar{\mathcal{U}}})$ is surjective; hence the map $K \to \bar{K}$ is also surjective. Q.E.D.

Remark (2.6). The results in this section equally hold in the complex analytic category. For example, let (X, p) be the germ of a symplectic variety X at $p \in X$, and let $f : (Y, E) \to (X, p)$ be a crepant, projective partial resolution of (X, p) where $E = f^{-1}(p)$. Assume that Y has only terminal singularities. Then (2.5) holds for (X, p) and (Y, E).

3 Symplectic automorphism and universal Poisson deformations

Let S be the same as in (1.2), and put $V := (S, 0) \times (\mathbf{C}^{2n-2}, 0)$. By the symplectic 2-form $\omega := (p_1)^*\omega_S + (p_2)^*\omega_{\mathbf{C}^{2n-2}}$, the germ V becomes a symplectic variety. Let $(\tilde{S}, F) \to (S, 0)$ be the minimal resolution and put $\tilde{V} := (\tilde{S}, F) \times (\mathbf{C}^{2n-2}, 0)$. In this section, we construct explicitly the universal Poisson deformations of V and $\tilde{V}$, and study the natural action on them induced by a symplectic automorphism of V. Let $\mathfrak{g}$ be the complex simple Lie algebra of the same type as S. Fix a Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$ and consider the adjoint quotient map $\mathfrak{g} \to \mathfrak{h}/W$, where W is the Weyl group of $\mathfrak{g}$. By [Slo], a transversal slice $\mathcal{S}$ of $\mathfrak{g}$ at the sub-regular nilpotent orbit gives the semi-universal flat deformation $\mathcal{S} \to \mathfrak{h}/W$ of S (at $0 \in \mathfrak{h}/W$). Let $\mathfrak{g}_{reg}$ be the open set of $\mathfrak{g}$ where this map is smooth. Then $\mathfrak{g}_{reg} \to \mathfrak{h}/W$ admits a relative symplectic 2-form called the Kostant-Kirillov 2-form. Let $\mathcal{S}_{reg}$ be the open subset of $\mathcal{S}$ where the map $\mathcal{S} \to \mathfrak{h}/W$ is smooth. The Kostant-Kirillov 2-form on $\mathfrak{g}_{reg}$ restricts to give a relative symplectic 2-form on $\mathcal{S}_{reg}$ and makes the map $\mathcal{S} \to \mathfrak{h}/W$ a Poisson deformation of S.

On the other hand, the base change $\mathfrak{g} \times_{\mathfrak{h}/W} \mathfrak{h} \to \mathfrak{h}$ has a simultaneous resolution

$$\mu : G \times^B \mathfrak{b} \to \mathfrak{g} \times_{\mathfrak{h}/W} \mathfrak{h},$$

where G is the adjoint group of $\mathfrak{g}$ and B is a Borel subgroup of G such that $\mathfrak{h} \subset \mathfrak{b}$ (cf. [Slo]). The pullback of the Kostant-Kirillov 2-form gives a relative symplectic 2-form $\omega_F \in \Gamma(G \times^B \mathfrak{b}, \Omega^2_{G\times^B\mathfrak{b}/\mathfrak{h}})$. If we put $\tilde{\mathcal{S}} := \mu^{-1}(\mathcal{S} \times_{\mathfrak{h}/W} \mathfrak{h})$, then

$$\mu|_{\tilde{\mathcal{S}}} : \tilde{\mathcal{S}} \to \mathcal{S} \times_{\mathfrak{h}/W} \mathfrak{h}$$

is a simultaneous resolution of $\mathcal{S} \times_{\mathfrak{h}/W} \mathfrak{h} \to \mathfrak{h}$. Let f be the composite of two maps $\tilde{\mathcal{S}} \to \mathcal{S} \times_{\mathfrak{h}/W} \mathfrak{h}$ and $\mathcal{S} \times_{\mathfrak{h}/W} \mathfrak{h} \to \mathfrak{h}$. Then $\omega_f := \omega_F|_{\tilde{\mathcal{S}}}$ gives a relative symplectic 2-form for f (cf. [Ya]).

Proposition (3.1) *(1) The universal Poisson deformations of S and $\tilde{S}$ are respectively given by $\mathcal{S} \to \mathfrak{h}/W$ and $\tilde{\mathcal{S}} \to \mathfrak{h}$.*

(2) The universal Poisson deformations of V and $\tilde{V}$ are respectively given by $\mathcal{S} \times (\mathbf{C}^{2n-2}, 0) \to \mathfrak{h}/W$ and $\tilde{\mathcal{S}} \times (\mathbf{C}^{2n-2}, 0) \to \mathfrak{h}$.

Proof. The Poisson deformation $\mathcal{S} \to \mathfrak{h}/W$ is universal at $0 \in \mathfrak{h}/W$. In fact, there is an exact sequence (cf. the latter part of §1 after (1.8))

$$0 \to \mathrm{PD}_{lt,S}(\mathbf{C}[\epsilon]) \to \mathrm{PD}_S(\mathbf{C}[\epsilon]) \to T^1_S \to 0.$$

For the definitions of PD and PD_{lt}, see (1.1) and (1.8). By Proposition (1.11), we have $\mathrm{PD}_{lt,S}(\mathbf{C}[\epsilon]) \cong H^2(S, \mathbf{C}) = 0$. The map $\mathrm{PD}_S(\mathbf{C}[\epsilon]) \to T^1_S$ is an isomorphism. Since $\mathcal{S} \to \mathfrak{h}/W$ is a semi-universal flat deformation of S, the Kodaira-Spencer map $T_{\mathfrak{h}/W,0} \to T^1_S$ is an isomorphism. The Kodaira-Spencer map factorizes as $T_{\mathfrak{h}/W,0} \to \mathrm{PD}_S(\mathbf{C}[\epsilon]) \to T^1_S$; hence the Poisson Kodaira-Spencer map $T_{\mathfrak{h}/W,0} \to \mathrm{PD}_S(\mathbf{C}[\epsilon])$ is an isomorphism. This fact together with (2.6) implies the universality of the Poisson deformation. Now let us consider the map $\tilde{\mathcal{S}} \to \mathfrak{h}$. By [Slo], it is semi-universal as a usual flat deformation of $\tilde{S}$. Therefore, the Kodaira-Spencer map $T_{\mathfrak{h},0} \to H^1(\tilde{S}, \Theta_{\tilde{S}})$ is an isomorphism. Moreover, this map factorizes as $T_{\mathfrak{h},0} \to H^2(\tilde{S}, \mathbf{C}) \to H^1(\tilde{S}, \Theta_{\tilde{S}})$, where the map $T_{\mathfrak{h},0} \to H^2(\tilde{S}, \mathbf{C})$ is the Poisson Kodaira-Spencer map. By the symplectic 2-form, $\Theta_{\tilde{S}}$ and $\Omega^1_{\tilde{S}}$ are identified. Then, the map $H^2(\tilde{S}, \mathbf{C}) \to H^1(\tilde{S}, \Theta_{\tilde{S}})$ coincides with the natural isomorphism $H^2(\tilde{S}, \mathbf{C}) \to H^1(\tilde{S}, \Omega^1_{\tilde{S}})$. Therefore, the Poisson Kodaira-Spencer map $T_{\mathfrak{h},0} \to H^2(\tilde{S}, \mathbf{C})$ is an isomorphism. This fact together with (2.6) implies that $f : \tilde{\mathcal{S}} \to \mathfrak{h}$ is the universal Poisson deformation of $\tilde{S}$. Let us now consider the Poisson deformations of $\tilde{V}$. The tangent space $\mathrm{PD}_{\tilde{V}}(\mathbf{C}[\epsilon])$ of the Poisson deformation functor is isomorphic to $H^2(\tilde{S} \times \mathbf{C}^{2n-2}, \mathbf{C}) = H^2(\tilde{S}, \mathbf{C})$. Since $\mathrm{PD}_{(\tilde{S},F)}(\mathbf{C}[\epsilon]) \cong H^2(\tilde{S}, \mathbf{C})$, this means that

$$\tilde{\mathcal{S}} \times \mathbf{C}^{2n-2} \overset{f\circ p_1}{\to} \mathfrak{h}$$

is the universal Poisson deformation of $\tilde{V}$ at $0 \in \mathfrak{h}$. Moreover, the map

$$\mathcal{S} \times \mathbf{C}^{2n-2} \to \mathfrak{h}/W$$

is the universal Poisson deformation of V at $0 \in \mathfrak{h}/W$. In fact, the map $\mathcal{S} \to \mathfrak{h}/W$ is the universal Poisson deformation of S. By Lemma (1.6), any 1-st order Poisson deformation is the product of a 1-st order Poisson deformation of S and $(\mathbf{C}^{2n-2}, 0)$. Then, the Poisson Kodaira-Spencer map $T_{\mathfrak{h}/W,0} \to \mathrm{PD}_V(\mathbf{C}[\epsilon])$ is an isomorphism. Q.E.D.

Let

$$i : V \to V$$

be a symplectic automorphism of V. The map i lifts to a symplectic automorphism

$$\tilde{i} : \tilde{V} \to \tilde{V}$$

so that the following diagram commutes

$$\begin{tikzcd} \tilde{V} \arrow[r, "\tilde{i}"] \arrow[d] & \tilde{V} \arrow[d] \\ V \arrow[r, "i"] & V. \end{tikzcd} \tag{2}$$

Correspondingly, we have a commutative diagram of functors:

$$\begin{tikzcd}
\mathrm{PD}_{\tilde{V}} \arrow[r, "\tilde{i}_*"] \arrow[d] & \mathrm{PD}_{\tilde{V}} \arrow[d] \\
\mathrm{PD}_V \arrow[r, "i"] & \mathrm{PD}_V .
\end{tikzcd} \tag{3}$$

By the (formal) universality of PD_V and $\mathrm{PD}_{\tilde{V}}$ (cf.(2.5), (2.6)), we have a commutative diagram

$$\begin{tikzcd}
\hat{\mathfrak{h}} \arrow[r, "\tilde{\iota}"] \arrow[d] & \hat{\mathfrak{h}} \arrow[d] \\
\mathfrak{h}/\hat{W} \arrow[r, "\iota"] & \mathfrak{h}/\hat{W},
\end{tikzcd} \tag{4}$$

where $\hat{\mathfrak{h}}$ and $\mathfrak{h}/\hat{W}$ are the formal completions of $\mathfrak{h}$ and $\mathfrak{h}/W$ at the origins.

Proposition (3.2) *The quotient space $\mathfrak{h}/W$ has a linear structure so that the commutative diagram above is obtained from a commutative diagram of linear spaces*

$$\begin{tikzcd}
\mathfrak{h} \arrow[r] \arrow[d] & \mathfrak{h} \arrow[d] \\
\mathfrak{h}/W \arrow[r] & \mathfrak{h}/W
\end{tikzcd} \tag{5}$$

where both horizontal maps are linear maps. Moreover, the horizontal map $\mathfrak{h} \to \mathfrak{h}$ is induced by a graph automorphism of the Dynkin diagram of $\mathfrak{g}$.

Proof. Let us consider the Poisson deformation $\tilde{\mathcal{S}} \times (\mathbf{C}^{2n-2}, 0) \to \mathfrak{h}$. The relative symplectic 2-form $\omega_f + \omega_{\mathbf{C}^{2n-2}}$ defines a 2-nd cohomology class of each fiber $\tilde{\mathcal{S}}_t \times (\mathbf{C}^{2n-2}, 0)$, $t \in \mathfrak{h}$. Since $H^2(\tilde{S}_t \times \mathbf{C}^{2n-2}, \mathbf{C})$ is identified with $H^2(\tilde{V}, \mathbf{C})$, one can define a period map (cf. [G-K], [Ya])

$$p : \mathfrak{h} \to H^2(\tilde{V}, \mathbf{C}) \cong H^2(\tilde{S}, \mathbf{C}).$$

Similarly one can define a period map

$$p_B : \mathfrak{h} \to H^2(T^*(G/B), \mathbf{C})$$

for the Poisson deformation $F : (G \times^B \mathfrak{b}) \times (\mathbf{C}^{2n-2}, 0) \to \mathfrak{h}$ by using the relative symplectic 2-form $\omega_F + \omega_{\mathbf{C}^{2n-2}}$. Since $\omega_f = \omega_F|_{\tilde{\mathcal{S}}}$, the period map p

is the composite of p_B and the natural restriction map $H^2(T^*(G/B), \mathbf{C}) \to H^2(\tilde{S}, \mathbf{C})$. This restriction map is an isomorphism since $\mathfrak{g}$ is simply-laced. Note that W has monodromy actions on $H^2(T^*(G/B), \mathbf{C})$ and $H^2(\tilde{S}, \mathbf{C})$ ([Slo 2], 4.2, 4.3, 4.4). By [Ya, Section 3] the period map p_B is a W-equivariant linear isomorphism; hence p is also a W-equivariant linear isomorphism. The description of p_B is as follows. First of all, the nilpotent cone N of $\mathfrak{g}$ is resolved by the Springer map $\mu_0 : T^*(G/B) \to N$. The transversal slice S is contained in N and $\tilde{S} = \mu_0^{-1}(S)$. There is an isomorphism

$$\mathfrak{h}^* \to H^2(T^*(G/B), \mathbf{C}).$$

The construction is as follows. Let $H \subset B$ be the maximal torus corresponding to $\mathfrak{h}$. Then there is a canonical isomorphism (cf. [Na 5, (P3)])

$$\mathrm{Hom}_{alg.gp}(H, \mathbf{C}^*) \otimes \mathbf{C} \cong \mathrm{Pic}(G/B) \otimes \mathbf{C}.$$

The left hand side is $\mathfrak{h}^*$ and right hand side is isomorphic to $H^2(G/B, \mathbf{C})$. Since $H^2(G/B, \mathbf{C}) \cong H^2(T^*(G/B), \mathbf{C})$, we have an isomorphism $\mathfrak{h}^* \to H^2(T^*(G/B), \mathbf{C})$. The Cartan subalgebra $\mathfrak{h}$ is identified with its dual $\mathfrak{h}^*$ by the Killing form of $\mathfrak{g}$. By §3 of [Ya] the period map p_B coincides with the composite of two maps:

$$\mathfrak{h} \to \mathfrak{h}^* \to H^2(T^*(G/B), \mathbf{C}).$$

The automorphism $\tilde{i}$ of $\tilde{V}$ induces an isomorphism

$$\tilde{i}^* : H^2(\tilde{V}, \mathbf{C}) \to H^2(\tilde{V}, \mathbf{C}).$$

By the identification $H^2(\tilde{V}, \mathbf{C}) \cong H^2(\tilde{S}, \mathbf{C})$, the map $\tilde{i}^*$ is regarded as an automorphism of $H^2(\tilde{S}, \mathbf{C})$. By the definition of $\tilde{\imath}$ we have a commutative diagram

$$\begin{tikzcd}
\hat{\mathfrak{h}} \arrow[r] & \mathfrak{h} \arrow[r, "p"] & H^2(\tilde{S}, \mathbf{C}) \arrow[d, "\tilde{i}^*"] \\
\hat{\mathfrak{h}} \arrow[u, "\tilde{\imath}"] \arrow[r] & \mathfrak{h} \arrow[r, "p"] & H^2(\tilde{S}, \mathbf{C}).
\end{tikzcd} \tag{6}$$

Define a linear map $\tilde{\imath}_{\mathfrak{h}} : \mathfrak{h} \to \mathfrak{h}$ by $p^{-1} \circ (\tilde{i}^*)^{-1} \circ p$. Then we have a commutative diagram

$$\begin{tikzcd}
\hat{\mathfrak{h}} \arrow[r] \arrow[d, "\tilde{\imath}"] & \mathfrak{h} \arrow[d, "\tilde{\imath}_{\mathfrak{h}}"] \\
\hat{\mathfrak{h}} \arrow[r] & \mathfrak{h}
\end{tikzcd} \tag{7}$$

We shall prove that $\tilde{\iota}_{\mathfrak{h}}$ is induced by a graph automorphism of $\mathfrak{g}$. Let $\Phi \subset \mathfrak{h}$ be the (co)root system for $\mathfrak{g}$. The choice of B determines a base Δ of Φ. Define

$$\Gamma := \{\phi \in \mathrm{Aut}(\Phi); \phi(\Delta) = \Delta\}.$$

Let C_i be a (-2)-curve on $\tilde{S}$ and let $[C_i] \in H^2(\tilde{S}, \mathbf{C})$ be its class. Define

$$\Phi' := \{C := \Sigma a_i[C_i]; a_i \in \mathbf{Z},\ C^2 = -2\}.$$

Then Φ' becomes a root system and $\Delta' := \{[C_i]\}$ forms a base of Φ'. Define

$$\Gamma' := \{\phi \in \mathrm{Aut}(\Phi'); \phi(\Delta') = \Delta'\}.$$

The period map p sends Δ to Δ' up to a non-zero constant. Since $\tilde{i}^* \in \Gamma'$, we have $\tilde{\iota}_{\mathfrak{h}} \in \Gamma$. The Weyl group W of $\mathfrak{g}$ is a normal subgroup of $\mathrm{Aut}(\Phi)$ and $\mathrm{Aut}(\Phi)$ is the semi-direct product of W and Γ. This means that $\tilde{\iota}_{\mathfrak{h}}$ descends to an automorphism $\iota_{\mathfrak{h}/W}$ of $\mathfrak{h}/W$. Since W is a finite reflection group, $\mathfrak{h}/W$ is an affine space. By [Slo, 8.8, Lemma 1], one can choose a linear structure of $\mathfrak{h}/W$ so that $\iota_{\mathfrak{h}/W}$ is a linear map.

4 Global sections of the local system

(4.1) **Monodromy of $R^2\pi^{an}_*\mathbf{C}$**

As in (1.2)-(1.5), we shall consider a symplectic variety (X, ω) whose singularities are locally isomorphic to $(S, 0) \times (\mathbf{C}^{2n-2}, 0)$. We use the same notation in section 1. Let $\pi : Y \to X$ be the minimal resolution. By definition, π^{an} is locally a product of the minimal resolution $\tilde{S} \to S$ and the $2n-2$ dimensional disc Δ^{2n-2}. If S is of type A_r, D_r or E_r, then, for each $p \in \Sigma$, the fiber $(\pi^{an})^{-1}(p)$ has r irreducible components and each of them is isomorphic to $\mathbf{P}^1$. Let E be the π-exceptional locus and let m be the number of irreducible components of E. We have $m \leq r$; but $m \neq r$ in general. The local system $R^2\pi^{an}_*\mathbf{C}$ on Σ may possibly have monodromies. Let γ be a closed loop in Σ starting from $p \in \Sigma$. Then we have a monodromy transformation along γ:

$$H^2((\pi^{an})^{-1}(p), \mathbf{C}) \to H^2((\pi^{an})^{-1}(p), \mathbf{C}).$$

Since $H^2((\pi^{an})^{-1}(p), \mathbf{C}) \cong H^2(\tilde{S}, \mathbf{C})$, the monodromy transformation is an automorphism of $H^2(\tilde{S}, \mathbf{C})$. Let F be the exceptional divisor of the minimal resolution $\tilde{S} \to S$ and let $F = \cup F_i$ be the irreducible decomposition.

Then $\{[F_i]\}$ is a basis of $H^2(\tilde{S}, \mathbf{C})$. The monodromy transformation permutes $[F_i]$'s without changing the intersection numbers. Therefore, the monodromy transformation comes from a graph automorphism of the Dynkin diagram associated with S. Let us observe the graph automorphisms of various Dynkin diagrams. In the (A_r)-case, the Dynkin diagram



has an automorphism σ_1 of order 2 which sends each i-th vertex to the $r+1-i$-th vertex. Hence, there are two possibilities for m; namely,

$$m = r, \text{ or } r - [r/2].$$

The Dynkin diagram of type D_r



has an automorphism σ_2 of order 2, which sends the 1-st vertex to the 2-nd one. Especially when $r = 4$, it has another automorphism τ of order 3 which permutes mutually the 1-st vertex, the 2-nd one and 3-rd one. Hence, in the (D_4)-case, there are three possibilities for m

$$m = 4, 3 \text{ or } 2,$$

and, in the (D_r)-case with $r > 4$, there are two possibilities for m

$$m = r \text{ or } r - 1.$$

Finally, let us consider the (E_6)-case.



The diagram has an automorphism σ_3 of order 2, which sends the 1-st vertex to the 6-th one and the 2-nd one to the 5-th one. There are two possibilities for m

$$m = 6, \text{ or } 4.$$

Since there are no symmetries for the diagrams of type (E_7), (E_8), we conclude that $m = r$ in these cases.

Let γ be a closed loop in Σ starting from $p \in \Sigma$. In (1.7), we have chosen a sequence of points p_i $(1 \leq i \leq k)$ on γ and have made a sequence of symplectic isomorphisms $(X^{an}, p_{i-1}) \cong (X^{an}, p_i)$. The composite of them finally defines a symplectic automorphism

$$i_\gamma : (X^{an}, p) \cong (X^{an}, p).$$

Here we shall describe the monodromy transformation of $R^2\pi^{an}_*\mathbf{C}$ along γ in terms of a symplectic automorphism of $(Y^{an}, \pi^{-1}(p))$. For each open set $V \subset X^{an}$, we associate the $\mathbf{C}$-vector space which consists of all 1-st order Poisson deformations of $(\pi^{an})^{-1}(V)$. The sheaf determined by this presheaf is isomorphic to $R^2\pi^{an}_*\mathbf{C}$ (cf. [Na 2]). The symplectic isomorphisms $(X^{an}, p_{i-1}) \cong (X^{an}, p_i)$ induce symplectic isomorphisms $(Y^{an}, (\pi^{an})^{-1}(p_{i-1})) \cong (Y^{an}, (\pi^{an})^{-1}(p_i))$ because $(Y^{an}, (\pi^{an})^{-1}(p_i))$ is a unique crepant resolution of (X^{an}, p_i). The sequence of them finally defines a symplectic automorphism

$$\tilde{i}_\gamma : (Y^{an}, (\pi^{an})^{-1}(p) \cong (Y^{an}, (\pi^{an})^{-1}(p)).$$

Note that $\tilde{i}_\gamma$ is a (unique) lift of i_γ to an automorphism of $(Y^{an}, (\pi^{an})^{-1}(p))$. The map $\tilde{i}_\gamma$ induces an automorphism of $(R^2\pi^{an}_*\mathbf{C})_p$, which is nothing but the monodromy transformation of $R^2\pi^{an}_*\mathbf{C}$ along γ. The identification $(X^{an}, p) \cong (S, 0) \times (\mathbf{C}^{2n-2}, 0)$ naturally lifts to the identification of $(Y^{an}, (\pi^{an})^{-1}(p))$ with $(\tilde{S}, F) \times (\mathbf{C}^{2n-2}, 0)$. Then, $(R^2\pi^{an}_*\mathbf{C})_p$ can be identified with $H^2(\tilde{S}, \mathbf{C})$.

The following is the main result in this section.

Proposition (4.2). *The following equality holds:*

$$\dim_{\mathbf{C}} H^0(\Sigma, \mathcal{H}) = m.$$

Proof. Let γ be a closed loop starting from $p \in \Sigma$. As in (1.7), we choose admissible covers $\{U_i\}$ of γ and points $p_i \in \Gamma$. By (1.7) and (4.1), the monodromy transformations of $\mathcal{H}_p$ and $(R^2\pi^{an}_*\mathbf{C})_p$ along γ, are described in terms of symplectic automorphisms

$$i_\gamma : (X^{an}, p) \to (X^{an}, p)$$

and

$$\tilde{i}_\gamma : (Y^{an}, (\pi^{an})^{-1}(p)) \to (Y^{an}, (\pi^{an})^{-1}(p)).$$

Apply Proposition (3.2) to these symplectic automorphisms. Then the sheaf $R^2\pi^{an}_*\mathbf{C}$ is a local system of the $\mathbf{C}$-module $\mathfrak{h}$, and $\mathcal{H}$ is a local system of the $\mathbf{C}$-module $\mathfrak{h}/W$. Moreover, their monodromies along γ are given by the horizontal maps $\mathfrak{h} \to \mathfrak{h}$ and $\mathfrak{h}/W \to \mathfrak{h}/W$ in the commutative diagram in Proposition (3.2). According to the notation in the proof of (3.2), we call these maps $\tilde{\iota}_{\gamma,\mathfrak{h}}$ and $\iota_{\gamma,\mathfrak{h}/W}$ respectively. Assume that S is of type A_r, D_r or E_r. When $m = r$, the sheaf $R^2\pi^{an}_*\mathbf{C}$ has a trivial monodromy along any γ. In this case, we have $\tilde{\iota}_{\gamma,\mathfrak{h}} = id$; hence $\iota_{\gamma,\mathfrak{h}/W} = id$. The problem is when $m < r$. In this case, there is a loop γ such that $\tilde{\iota}_{\gamma,\mathfrak{h}}$ comes from one of the graph automorphisms listed in (4.1). Assume that $\dim \mathfrak{h}^{\tilde{\iota}_{\gamma,\mathfrak{h}}} = m$, where $\mathfrak{h}^{\tilde{\iota}_{\gamma,\mathfrak{h}}}$ is the invariant part of $\mathfrak{h}$ under $\tilde{\iota}_{\gamma,\mathfrak{h}}$. By the argument in [Slo, 8.8, Lemma 1], we see that $\dim(\mathfrak{h}/W)^{\iota_{\gamma,\mathfrak{h}/W}} = m$. Q.E.D.

By using Proposition (4.2), we can prove that the inequality in Corollary (1.10) of [Na 1] is actually an equality:

Corollary (4.3). *Let (X, ω) be a projective symplectic variety. Let $U \subset X$ be the locus where X is locally a trivial deformation of a (surface) rational double point at each $p \in U$. Let $\pi : \tilde{U} \to U$ be the minimal resolution and let m be the number of irreducible components of* $\mathrm{Exc}(\pi)$. *Then $h^0(U, T^1_U) = m$.*

Proof By Lemma (1.5) we obtain a local system $\mathcal{H}$ of $\mathbf{C}$-modules as a subsheaf of T^1_U. Put $\Sigma := \mathrm{Sing}(U)$. Let $\Sigma = \cup\Sigma_i$ be the decomposition into connected components. The local system $\mathcal{H}$ has support on Σ. Let $\mathcal{H}_i$ be the restriction of $\mathcal{H}$ to each connected component Σ_i. We have an isomorphism:

$$\mathcal{H} \otimes_{\mathbf{C}} \mathcal{O}_\Sigma \cong T^1_U.$$

Then

$$h^0(U, T^1_U) = h^0(\Sigma, \mathcal{H} \otimes_{\mathbf{C}} \mathcal{O}_\Sigma) = \Sigma h^0(\mathcal{H}_i) \cdot h^0(\mathcal{O}_{\Sigma_i}).$$

Since Σ_i can be compactified to a proper normal variety $\bar{\Sigma}_i$ such that $\bar{\Sigma}_i - \Sigma_i$ has codimension ≥ 2, we see that $h^0(\mathcal{O}_{\Sigma_i}) = 1$. Q.E.D.

5 Main Results

Theorem (5.1). *Let X be an affine symplectic variety. Then* PD_X *is unobstructed.*

Proof. (i) Let U be chosen as in (1.8). Let $\pi : \tilde{U} \to U$ be the minimal resolution. Put $Z := X \setminus U$. In the exact sequence of local cohomology

$$... \to H^i(X, \mathcal{O}_X) \to H^i(U, \mathcal{O}_U) \to H^{i+1}_Z(X, \mathcal{O}_X) \to ...,$$

we have $H^{i+1}_Z(X, \mathcal{O}_X) = 0$ for all $i \leq 2$ since X is Cohen-Macaulay and $\mathrm{Codim}_X Z \geq 4$. Note that $H^i(X, \mathcal{O}_X) = 0$ for $i > 0$. Therefore, one has $H^i(U, \mathcal{O}_U) = 0$ for $i = 1, 2$. Since U is a symplectic variety, U has only rational singularities (cf. (1.1)). In particular, this implies that $H^i(\tilde{U}, \mathcal{O}_{\tilde{U}}) = 0$ for $i = 1, 2$. The resolution $\tilde{U}$ is a smooth symplectic variety and $\mathrm{PD}_{\tilde{U}}(\mathbf{C}[\epsilon]) \cong H^2(\tilde{U}^{an}, \mathbf{C})$. There is a natural map $\mathrm{PD}_{\tilde{U}}(\mathbf{C}[\epsilon]) \to \mathrm{PD}_U(\mathbf{C}[\epsilon])$. In fact, since $R^1\pi_*\mathcal{O}_{\tilde{U}} = 0$ and $\pi_*\mathcal{O}_{\tilde{U}} = \mathcal{O}_U$, a first order deformation $\tilde{\mathcal{U}}$ (without Poisson structure) of $\tilde{U}$ induces a first order deformation $\mathcal{U}$ of U (cf. [Wa]). Let $\mathcal{U}^0$ be the locus where $\mathcal{U} \to \mathrm{Spec}(\mathbf{C}[\epsilon])$ is smooth. Since $\tilde{\mathcal{U}} \to \mathcal{U}$ is an isomorphism above $\mathcal{U}^0$, the Poisson structure of $\tilde{\mathcal{U}}$ induces that of $\mathcal{U}^0$. Since the Poisson structure of $\mathcal{U}^0$ uniquely extends to that of $\mathcal{U}$, $\mathcal{U}$ becomes a Poisson scheme over $\mathrm{Spec}(\mathbf{C}[\epsilon])$. This is the desired map. In the same way, one has a morphism of functors:

$$\mathrm{PD}_{\tilde{U}} \overset{\pi_*}{\to} \mathrm{PD}_U.$$

Note that $\mathrm{PD}_{\tilde{U}}$ (resp. PD_U) has a prorepresentable hull $R_{\tilde{U}}$ (resp. R_U). Then π_* induces a local homomorphism of complete local rings:

$$R_U \to R_{\tilde{U}}.$$

We now obtain a commutative diagram of exact sequences:

$$\begin{tikzcd}
0 \arrow[r] & H^2(U^{an}, \mathbf{C}) \arrow[r] \arrow[d, "\cong"'] & \mathrm{PD}_{\tilde{U}}(\mathbf{C}[\epsilon]) \arrow[r] \arrow[d] & H^0(U^{an}, R^2\pi^{an}_*\mathbf{C}) \\
0 \arrow[r] & \mathrm{PD}_{lt,U}(\mathbf{C}[\epsilon]) \arrow[r] & \mathrm{PD}_U(\mathbf{C}[\epsilon]) \arrow[r] & H^0(\Sigma, \mathcal{H})
\end{tikzcd} \quad (8)$$

(ii) Let E_i ($i = 1, ..., m$) be the irreducible components of $\mathrm{Exc}(\pi)$. Each E_i defines a class $[E_i] \in H^0(U^{an}, R^2\pi^{an}_*\mathbf{C})$. It is easily checked that $H^0(U^{an}, R^2\pi^{an}_*\mathbf{C}) = \oplus_{1 \leq i \leq m}\mathbf{C}[E_i]$. This means that

$$\dim \mathrm{PD}_{\tilde{U}}(\mathbf{C}[\epsilon]) = h^2(U^{an}, \mathbf{C}) + m.$$

On the other hand, by Proposition (4.2), $h^0(\Sigma, \mathcal{H}) = m$. This means that

$$\dim \mathrm{PD}_U(\mathbf{C}[\epsilon]) \leq h^2(U^{an}, \mathbf{C}) + m.$$

As a consequence, we have

$$\dim \mathrm{PD}_U(\mathbf{C}[\epsilon]) \leq \dim \mathrm{PD}_{\tilde{U}}(\mathbf{C}[\epsilon]).$$

(iii) We shall prove that the morphism $\pi_* : \mathrm{PD}_{\tilde{U}} \to \mathrm{PD}_U$ has a finite fiber. More exactly, $\mathrm{Spec}(R_{\tilde{U}}) \to \mathrm{Spec}(R_U)$ has a finite closed fiber. Let $\alpha : R_{\tilde{U}} \to \mathbf{C}[[t]]$ be a homomorphism of local $\mathbf{C}$-algebras such that the composition map $R_U \to R_{\tilde{U}} \overset{\alpha}{\to} \mathbf{C}[[t]]$ is factorized as $R_U \to R_U/m_U \to \mathbf{C}[[t]]$. Let U_p be the germ of U^{an} at $p \in \Sigma$ and let $\tilde{U}_p$ be the germ of $\tilde{U}^{an}$ along $(\pi^{an})^{-1}(p)$. Denote by R_{U_p} (resp. $R_{\tilde{U}_p}$) the prorepresentable hull of the Poisson deformation functor PD_{U_p} (resp. $\mathrm{PD}_{\tilde{U}_p}$). Since a Poisson deformation of U (resp. $\tilde{U}$) induces a Poisson deformation of U_p (resp. $\tilde{U}_p$), α induces the map $\alpha_p : R_{\tilde{U}_p} \to \mathbf{C}[[t]]$ such that the map $R_{U_p} \to R_{\tilde{U}_p} \overset{\alpha_p}{\to} \mathbf{C}[[t]]$ is factorizes as $R_{U_p} \to R_{U_p}/m_{U_p} \to \mathbf{C}[[t]]$. Corresponding to α, we have a family of morphisms $\{\pi_n\}_{n\geq 1}$:

$$\pi_n : \tilde{U}_n \to U_n,$$

where $U_n \cong U \times \mathrm{Spec}\mathbf{C}[t](t^{n+1})$ and $\tilde{U}_n$ are Poisson deformations of $\tilde{U}$ over $\mathrm{Spec}\mathbf{C}[t]/(t^{n+1})$. Restrict these to $\tilde{U}_p$ and U_p. Then we have a family of morphisms $\{\pi_{p,n}\}_{n\geq 1}$:

$$\pi_{p,n} : \tilde{U}_{p,n} \to U_{p,n},$$

which are Poisson deformations of $\tilde{U}_p$ and U_p determined by α_p. As proved in (3.1), the map $\mathrm{Spec}(R_{\tilde{U}_p}) \to \mathrm{Spec}(R_{U_p})$ is a finite Galois covering. This means that each $\tilde{U}_{p,n}$ coincides with the minimal resolution of $U_{p,n}$ (i.e. $\tilde{U}_p \times \mathrm{Spec}\mathbf{C}[t]/(t^{n+1})$) with the natural Poisson structure determined by that of $U_{p,n}$. Since all minimal resolution $\tilde{U}_{p,n}$ ($p \in \Sigma$) are glued together, we conclude that $\tilde{U}_n \cong \tilde{U} \times \mathrm{Spec}\mathbf{C}[t]/(t^{n+1})$ and its Poisson structures is uniquely determined by that of U_n. This implies that the given map $R_{\tilde{U}} \to \mathbf{C}[[t]]$ factors through $R_{\tilde{U}}/m_{\tilde{U}}$.

(iv) Since the tangent space of $\mathrm{PD}_{\tilde{U}}$ is controlled by $H^2(U^{an}, \mathbf{C})$, it has the T^1-lifting property; hence $\mathrm{PD}_{\tilde{U}}$ is unobstructed and $R_{\tilde{U}}$ is regular.

(v) By (ii), (iii) and (iv), we conclude that R_U is a regular local ring with $\dim R_U = \dim R_{\tilde{U}}$. In fact, since $\dim R_{\tilde{U}} \leq \dim R_U + \dim R_{\tilde{U}}/m_U R_{\tilde{U}}$, we have

$$\dim R_{\tilde{U}} \leq \dim R_U$$

by (iii). Since $R_{\tilde{U}}$ is regular by (iv), we have an equality

$$\dim_{\mathbf{C}} m_{\tilde{U}}/(m_{\tilde{U}})^2 = \dim R_{\tilde{U}}.$$

On the other hand, we have an inequality

$$\dim_{\mathbf{C}} m_U/(m_U)^2 \geq \dim R_U.$$

These three (in)equalities imply that

$$\dim_{\mathbf{C}} m_U/(m_U)^2 \geq \dim_{\mathbf{C}} m_{\tilde{U}}/(m_{\tilde{U}})^2.$$

Finally, by (ii), we see that this inequality actually is an equality, and the equality $\dim R_U = \dim_{\mathbf{C}} m_U/(m_U)^2$ holds.

Moreover, in the commutative diagram above, the map $\mathrm{PD}_U(\mathbf{C}[\epsilon]) \to H^0(\Sigma, \mathcal{H})$ is surjective. We shall prove that PD_X is unobstructed. Let $S_n := \mathbf{C}[t]/(t^{n+1})$ and $S_n[\epsilon] := \mathbf{C}[t, \epsilon]/(t^{n+1}, \epsilon^2)$. Put $T_n := \mathrm{Spec}(S_n)$ and $T_n[\epsilon] := \mathrm{Spec}(S_n[\epsilon])$. Let X_n be a Poisson deformation of X over T_n. Define $\mathrm{PD}(X_n/T_n, T_n[\epsilon])$ to be the set of equivalence classes of the Poisson deformations of X_n over $T_n[\epsilon]$. The X_n induces a Poisson deformation U_n of U over T_n. Define $\mathrm{PD}(U_n/T_n, T_n[\epsilon])$ in a similar way. Then, by the same argument as [Na 2, Proposition 13], we have

$$\mathrm{PD}(X_n/T_n, T_n[\epsilon]) \cong \mathrm{PD}(U_n/T_n, T_n[\epsilon]).$$

Now, since PD_U is unobstructed, PD_U has the T^1-lifting property. This equality shows that PD_X also has the T^1-lifting property. Therefore, PD_X is unobstructed. Q.E.D.

(5.2) Let X be an affine symplectic variety. Take a (projective) resolution $Z \to X$. By Birkar-Cascini-Hacon-McKernan [B-C-H-M], one applies the minimal model program to this morphism and obtains a relatively minimal model $\pi : Y \to X$. The following properties are satisfied:

(i) π is a crepant, birational projective morphism.
(ii) Y has only $\mathbf{Q}$-factorial terminal singularities.

Note that Y naturally becomes a symplectic variety. Let $U \subset X$ be the open locus where, for each $p \in U$, the germ (X, p) is non-singular or the product of a surface rational double point and a non-singular variety. We put $\tilde{U} := \pi^{-1}(U)$. As in (i) of the proof of Theorem (5.1), the birational maps

π and $\pi|_{\tilde{U}}$ induces natural maps of functors $\pi_* : \mathrm{PD}_Y \to \mathrm{PD}_X$ and $(\pi|_{\tilde{U}})_* : \mathrm{PD}_{\tilde{U}} \to \mathrm{PD}_U$. There is a commutative diagram of Poisson deformation functors

$$\begin{tikzcd}
\mathrm{PD}_Y \arrow[r] \arrow[d] & \mathrm{PD}_{\tilde{U}} \arrow[d] \\
\mathrm{PD}_X \arrow[r] & \mathrm{PD}_U
\end{tikzcd} \tag{9}$$

and correspondingly a commutative diagram of prorepresentable hulls

$$\begin{tikzcd}
R_{\tilde{U}} \arrow[r] & R_Y \\
R_U \arrow[r] \arrow[u] & R_X \arrow[u]
\end{tikzcd} \tag{10}$$

Lemma (5.3). *The horizontal maps $R_{\tilde{U}} \to R_Y$ and $R_U \to R_X$ are both isomorphisms.*

Proof. Let V be the regular locus of Y. Then $\tilde{U}$ is contained in V, and we have the restriction map $H^2(V^{an}, \mathbf{C}) \to H^2(\tilde{U}^{an}, \mathbf{C})$. This map is an isomorphism by the proof of [Na 3], Proposition 2. Note that $\mathrm{PD}_Y(\mathbf{C}[\epsilon]) = H^2(V^{an}, \mathbf{C})$ and $\mathrm{PD}_{\tilde{U}}(\mathbf{C}[\epsilon]) = H^2(\tilde{U}, \mathbf{C})$. By the T^1-lifting principle, PD_Y and $\mathrm{PD}_{\tilde{U}}$ are both unobstructed. Let us consider the map $R_{\tilde{U}} \to R_Y$. By the observation above, $R_{\tilde{U}}$ and R_Y are both regular and the map induces an isomorphism of Zariski tangent spaces; hence $R_{\tilde{U}} \cong R_Y$. Next let us consider the map $R_U \to R_X$. By Theorem (5.1), both local rings are regular and the map induces an isomorphism of Zariski tangent spaces; hence $R_U \cong R_X$. Q.E.D.

By Theorem (5.1), $\dim R_U = \dim R_{\tilde{U}}$ and the closed fiber of $R_U \to R_{\tilde{U}}$ is finite; hence $\dim R_X = \dim R_Y$ and the closed fiber of $\pi^* : R_X \to R_Y$ is finite. By the generalized Weierstrass preparation theorem, R_Y is a finite R_X-module; in other words, $\mathrm{Spec}\, R_Y \to \mathrm{Spec}\, R_X$ is a finite morphism.

We put $R_{X,n} := R_X/m^n$ and $R_{Y,n} := R_Y/(m_Y)^n$. Since PD_X and PD_Y are both prorepresentable, there is a commutative diagram of formal universal deformations of X and Y:

$$\begin{tikzcd}
\{Y_n\}_{n\geq 1} \arrow[r] \arrow[d] & \{X_n\}_{n\geq 1} \arrow[d] \\
\mathrm{Spec}(R_{Y,n}) \arrow[r] & \mathrm{Spec}(R_{X,n})
\end{tikzcd} \tag{11}$$

(5.4) **Algebraization**

Let us assume that an affine symplectic variety (X, ω) satisfies the following condition (*).

(*)

(1) There is a $\mathbf{C}^*$-action on X with only positive weights and a unique fixed point $0 \in X$.

(2) The symplectic form ω has positive weight $l > 0$.

By Step 1 of Proposition (A.7) in [Na 2], the $\mathbf{C}^*$-action on X uniquely extends to the action on Y. These $\mathbf{C}^*$-actions induce those on R_X and R_Y. By Section 4 of [Na 2], R_Y is isomorphic to the formal power series ring $\mathbf{C}[[y_1, ..., y_d]]$ with $wt(y_i) = l$. Since $R_X \subset R_Y$, the $\mathbf{C}^*$-action on R_X also has positive weights. We put $A := \underleftarrow{\lim}\Gamma(X_n, \mathcal{O}_{X_n})$ and $B := \underleftarrow{\lim}\Gamma(Y_n, \mathcal{O}_{Y_n})$. Let $\hat{A}$ and $\hat{B}$ be the completions of A and B along their maximal ideals. Then one has the commutative diagram

$$\begin{tikzcd}
R_X \arrow[r] \arrow[d] & R_Y \arrow[d] \\
\hat{A} \arrow[r] & \hat{B}
\end{tikzcd} \tag{12}$$

Let S (resp. T) be the $\mathbf{C}$-subalgebra of $\hat{A}$ (resp. $\hat{B}$) generated by the eigen-vectors of the $\mathbf{C}^*$-action. On the other hand, the $\mathbf{C}$-subalgebra of R_Y generated by eigen-vectors, is nothing but $\mathbf{C}[y_1, ..., y_d]$. Let us consider the $\mathbf{C}$-subalgebra of R_X generated by eigen-vectors. By [Na 2], Lemma (A.2), it is generated by eigenvectors that form a basis of $m_X/(m_X)^2$. Since R_X is regular of the same dimension as R_Y, the subalgebra is a polynomial ring $\mathbf{C}[x_1, ..., x_d]$. Now the following commutative diagram algebraizes the previous diagram:

$$\begin{tikzcd}
\mathbf{C}[x_1, ..., x_d] \arrow[r] \arrow[d] & \mathbf{C}[y_1, ..., y_d] \arrow[d] \\
S \arrow[r] & T
\end{tikzcd} \tag{13}$$

By Theorem (5.4.5) of [EGA III], the (formal) birational projective morphism

$$Y_n \to \mathrm{Spec}(\hat{B}/(m_{\hat{B}})^n)$$

is algebraized to a birational projective morphism

$$\hat{Y} \to \mathrm{Spec}(\hat{B}).$$

Moreover, by a method similar to that in Appendix of [Na 2], this is further algebraized to

$$\mathcal{Y} \to \mathrm{Spec}(T).$$

If we put $\mathcal{X} := \mathrm{Spec}(S)$, then we have a $\mathbf{C}^*$-equivariant commutative diagram of algebraic schemes

$$\begin{tikzcd}
\mathcal{Y} \arrow[r] \arrow[d] & \mathcal{X} \arrow[d] \\
\mathrm{Spec}\mathbf{C}[y_1, ..., y_d] \arrow[r, "\psi"] & \mathrm{Spec}\mathbf{C}[x_1, ..., x_d]
\end{tikzcd} \tag{14}$$

Theorem (5.5). *In the diagram above,*

(a) *the map ψ is a finite surjective map,*

(b) $\mathcal{Y} \to \mathrm{Spec}\ \mathbf{C}[y_1, ..., y_d]$ *is a locally trivial deformation of Y, and*

(c) *the induced birational map $\mathcal{Y}_t \to \mathcal{X}_{\psi(t)}$ is an isomorphism for a general $t \in \mathrm{Spec}\mathbf{C}[y_1, ..., y_d]$.*

Proof. (a) follows from [Na 2], Lemma (A.4) since R_Y is a R_X-finite module.

(b): Since Y is $\mathbf{Q}$-factorial, Y^{an} is also $\mathbf{Q}$-factorial by Proposition (A.9) of [Na 2]. Then (b) is Theorem 17 of [Na 2].

(c) follows from Proposition 24 of [Na 2].

Corollary (5.6). *Let (X, ω) be an affine symplectic variety with the property (*). Then the following two conditions are equivalent:*

(1) *X has a crepant projective resolution.*

(2) *X has a smoothing by a Poisson deformation.*

Proof. (1) ⇒ (2): If X has a crepant resolution, say Y. By using this Y, one can construct a diagram in Theorem (5.5). Then, by the property (c), we see that X has a smoothing by a Poisson deformation.

(2) ⇒ (1): Let Y be a crepant $\mathbf{Q}$-factorial terminalization of X. It suffices to prove that Y is smooth. We again consider the diagram in Theorem (5.5). By the assumption, $\mathcal{X}_s$ is smooth for a general point $s \in \mathrm{Spec}\mathbf{C}[x_1, ..., x_d]$. By the property (a), one can find $t \in \mathrm{Spec}\mathbf{C}[y_1, ..., y_d]$ such that $\psi(t) = s$. By (c), one has an isomorphism $\mathcal{Y}_t \cong \mathcal{X}_s$. In particular, $\mathcal{Y}_t$ is smooth. Then, by (b), $Y (= \mathcal{Y}_0)$ is smooth.

References

[Ar] Artin, M.: On the solutions of analytic equations, Invent. Math. **5** 277-291 (1968)

[A-M] Atiyah, M., MacDonald, I.: Introduction to commutative algebra, Addison-Wesley

[Be] Beauville, A.: Symplectic singularities, Invent. Math. **139** (2000), 541-549

[BCHM] Birkar, C., Cascini, P., Hacon, C., McKernan, J.: Existence of minimal models for varieties of log general type, math.AG/0610203 (to appear in J. Amer. Math. Soc.)

[C-M] Collingwood, D. , McGovern, W. : Nilpotent orbits in semi-simple Lie algebras, van Nostrand Reinhold, Math. Series, 1993

[EGA III] Grothendieck, A.: Elements de geometrie algebraique III. Etude cohomologique des faisceaux coherents I, Publ.IHES **11** (1961), 167 pp

[Fu] Fu, B. : Symplectic resolutions for nilpotent orbits, Invent. Math. **151**. (2003), 167-186

[Fu 2] Fu, B.: On **Q**-factorial terminalizations of nilpotent orbits, math/0809.5109, to appear in J. Math. Pures Appl.

[G-K] Ginzburg, V., Kaledin, D.: Poisson deformations of symplectic quotient singularities, Adv. Math. **186** (2004), 1-57

[Gr] Grothendieck, A.: On the De Rham cohomology of algebraic varieties, Publ. Math. IHES **29** (1966), 95-103

[Ka 1] Kaledin, D.: Symplectic singularities from the Poisson view point, J. Reine Angew. Math. **600** (2006), 135-156

[Ka 2] Kaledin, D.: Symplectic resolutions: deformations and birational maps, math.AG/0012008

[Mo] Moser, J.: On the volume elements on a manifold. Trans. Amer. Math. Soc. **120** (1965), 286-294

[Na 1] Namikawa, Y.: Deformation theory of singular symplectic n-folds, Math. Ann. **319** (2001) 597-623

[Na 2] Namikawa, Y.: Flops and Poisson deformations of symplectic varieties, Publ. RIMS, Kyoto Univ. **44**, no. 2 (2008) 259-314

[Na 3] Namikawa, Y.: On deformations of $\mathbf{Q}$-factorial symplectic varieties, J. Reine Angew. Math. **599** (2006) 97-110

[Na 4] Namikawa, Y.: Induced nilpotent orbits and birational geometry, math.arxiv: 0809.2320

[Na 5] Namikawa, Y.: Birational geometry and deformations of nilpotent orbits, Duke Math. J. **143** (2008) 375-405

[Sch] Schlessinger, M.: Functors of Artin rings, Trans. Amer. Math. Soc. **130** (1968), 208-222

[Slo] Slodowy, P. : Simple singularities and simple algebraic groups. Lect. Note Math. **815**, Springer-Verlag, 1980

[Slo 2] Slodowy, P.: Four lectures on simple groups and singularities, Comm. of Math. Inst. Rijksuniversiteit Utrecht 1980

[St] Steenbrink, J.: Mixed Hodge structure on the vanishing cohomology. Real and complex singularities (Proc. Ninth Nordic Summer School/NAVF Sympos. Math., Oslo, 1976), pp. 525–563. Sijthoff and Noordhoff, Alphen aan den Rijn, 1977

[Wa] Wahl, J.: Equisingular deformations of normal surface singularities I, Ann. of Math. (2) **104** (1976), no. 2, 325-356.

[Wav] Wavrik, J.: Obstructions to the existence of a space of moduli, in Global Analysis, papers in honor of K. Kodaira, edited by D.C. Spencer and S. Iyanaga, Uni. of Tokyo Press and Princeton Univ. Press (1969), 403-414

[Ya] Yamada, H.: Lie group theoretical construction of period mapping, Math. Z. **220** (1995), 231-255

Yoshinori Namikawa
Department of Mathematics, Faculty of Science, Kyoto University, JAPAN
namikawa@math.kyoto-u.ac.jp